

12-1-2011

Section: Botany, Microbiology and Zoology

REPRODUCTIVE BIOLOGY OF THE FLATTENED PORCELAIN CRAB, PETROLISTHES RUFESCENS (PORCELLANIDAE: ANOMURA: CRUSTACEA) FROM AIN SUKHNA, GULF OF SUEZ, EGYPT

AWAAD EL-SAYED

Section of Marine Biology and Fishes, Zoology Department, Faculty of Science, Al- Azhar University, Cairo, Egypt.

MOSTAFA FOU DA

Consultant for Minister of Environment, State Ministry of Environment, Cairo, Egypt.

AHMED AZAB

Section of Marine Biology and Fishes, Zoology Department, Faculty of Science, Al- Azhar University, Cairo, Egypt.

ISMAIL ISMAIL

M.Sc. in Marine Biology.

Follow this and additional works at: <https://absb.researchcommons.org/journal>



Part of the [Life Sciences Commons](#)

How to Cite This Article

EL-SAYED, AWAAD; FOU DA, MOSTAFA; AZAB, AHMED; and ISMAIL, ISMAIL (2011) "REPRODUCTIVE BIOLOGY OF THE FLATTENED PORCELAIN CRAB, PETROLISTHES RUFESCENS (PORCELLANIDAE: ANOMURA: CRUSTACEA) FROM AIN SUKHNA, GULF OF SUEZ, EGYPT," *Al-Azhar Bulletin of Science*: Vol. 22: Iss. 2, Article 6.

DOI: <https://doi.org/10.21608/absb.2011.7030>

This Original Article is brought to you for free and open access by Al-Azhar Bulletin of Science. It has been accepted for inclusion in Al-Azhar Bulletin of Science by an authorized editor of Al-Azhar Bulletin of Science. For more information, please contact kh_Mekheimer@azhar.edu.eg.

REPRODUCTIVE BIOLOGY OF THE FLATTENED PORCELAIN CRAB, *PETROLISTHES RUFESCENS* (PORCELLANIDAE: ANOMURA: CRUSTACEA) FROM AIN SUKHNA, GULF OF SUEZ, EGYPT

AWAAD ABDU MOHAMMED EL-SAYED ^(*); MOSTAFA MOKHTAR FOUDA ^(**);
AHMED MOSAD AZAB ^(*) and ISMAIL MASOUD ISMAIL ^(***)

* *Section of Marine Biology and Fishes, Zoology Department, Faculty of Science, Al- Azhar University, Cairo, Egypt.*

** *Consultant for Minister of Environment, State Ministry of Environment, Cairo, Egypt.*

*** *M.Sc. in Marine Biology.*

Abstract

The reproductive biology of the porcelain crab *Petrolisthes rufescens* (Heller, 1861) population at Ain Sukhna (62 km south Suez City) was studied during the period from April 2000 to May 2001. Sexual dimorphism was determined based on morphological differences between sexes. Abdomen is semi-circular with four pairs of biramous appendages in females, but tapering with only two pairs of unequal uniramous appendages (pleopods) in males. Size at first maturity was determined using allometric growth occurring at puberty from the relationships between male's 2nd pleopodal length and chelae length against carapace length (C.L.) and by the appearance of distinctive ovaries in females. Overall sex ratio for this crab species was 1: 0.70 males to females, with highly significant Chi-square value ($X^2 = 7.486$, $P < 0.05$, $df = 1$). It was not constant throughout the year particularly during breeding seasons, showing highly significant Chi-square value ($X^2 = 20.185$, $df = 25$, $P < 0.05$). Based on the appearance of ovigerous females, the breeding of *P. rufescens* is lengthy, start often at late winter and early spring and extends through summer to early autumn. Incubated eggs are oval or semi-oval, sometimes elliptical with polygonal surfaces, and varied in size from 400 to 1000 μm with an average of $640 \pm 105 \mu\text{m}$. These eggs are passed through five stages of maturation, and showed remarkably gradual decreasing in size throughout the period of incubation, reaching the minimum at last stages (stage V or the releasing larvae). The color of incubated eggs is also changed from bright yellow in newly laid eggs to grey at the last stage of development. The fecundity of ovigerous females was relatively low, varied from 35 to 450 with an average of 208 ± 92 eggs /female crab. It was correlated by a curvilinear relationship with carapace length only.

Introduction

Porcelain crabs are much alike true crabs (brachyurans) but they can easily distinguished from them by possessing uropods and their small, dorsally directed 5th pair (last walking pairs) of legs under the carapace (Barnard, 1950; Fransozo and Bertini, 2001; Osawa and McLaughlin, 2010). These crabs belong to family Porcellanidae Haworth, 1825 (Anomura: Decapoda), which comprises about 27 to 30 genera including from 225 to 277 species (Melo, 1999; Fransozo and Bertini, 2001; Osawa and McLaughlin, 2010), most of them are worldwide in distribution, occupying a variety of littoral habitats living in hard substrates within crevices or under boulders and even in bottom covered by calcareous algae, excepting rare species living in the deep sea (Veloso and Melo, 1993).

Among Porcellanidae, genus *Petrolisthes* is the largest, accommodates about 102 species of the total belonging to this family, several of them are worldwide in their distribution (Haig, 1960; Hollebone and Hay, 2007; Ng *et al.*, 2008; McLaughlin *et*

al., 2007). In the Red Sea and its associated gulfs (Suez and Aqaba), genus *Petrolisthes* is represented by 8 valid species (Lewinsohn, 1969), most of them are found in the intertidal zones, but few extend into about 5 m depth in the shallow subtidal areas (Lewinsohn, 1969; El-Sayed, 2002; Fouda *et al.*, 2003).

At the northwestern coast of the Suez Gulf, particularly Ain Sukhna, the porcelain crab, *Petrolisthes rufescens* (Heller, 1861), is a common intertidal species. It inhabits, with the co-existing less common porcelain species, *Petrolisthes leptochelae* (Heller, 1861), the rocky and mixed (rocky and sandy) shores. During low tides, the individuals of this species escape always under small stones and boulders to protect itself from the effect of heat, desiccation and terrestrial predators (Ismail, 2005). They contribute with a significant part in the composition of the intertidal marine fauna at that area and can be used as a biological indicator for the effects of human activities on marine environment.

Since the last decade of the 20th century till now, several resorts, tourist villages, hotels and camp sites are either constructed or are under constructions along nearly every space at the western coastal-line of the Gulf of Suez extending from Suez City (north) to Za'afarana (south). Hence, the natural habitats at several intertidal areas became threatened as results of severe cleaning, dredges, filling and continuous changes for the natural profile of the coast-line, in addition to the profound effects of oil refiners and new constructed industrial commercial companies at Ain Sukhna. These man-made constructions associated with processes of reconstruction for the major areas of the shoreline and human activities, threat in turn and affect directly different types of littoral marine habitat with its associated fauna, particularly those distributed on the exposed intertidal and coastal zones. Therefore, particular attention needs to carry out detailed studies at that area, to be considered as baseline data available to follow up these changes and take in consideration the future studies which will be done later.

Consequently, the present work aims to through light on the reproductive biology of the flattened porcelain crab, *Petrolisthes rufescens*, one of the common species inhabiting the intertidal zones at the study area, due to its sensitivity to environmental changes and its important ecological role in food chains for other co-existing species.

Materials and Methods

The specimens of porcelain crab, *Petrolithes rufescens* were collected, from the intertidal and uppermost borders of subtidal zones of the five selected study areas along the northwestern portion of the Gulf of Suez (Figure 1) during the period from June 2000 to May 2001. The intensive monthly work and specimen collection were carried out at site # 1, which is known as Al- Fangary Beach, located about 2 km south Ain Sukhna (Km 62 South Suez City). Additional specimens were collected irregularly during the same period from the other four localities including: Al-Khaima Beach, site # 2 (40 km South Suez City), Al-Semad Beach, site # 3 (32 km South Suez City), the shore line of the National Institute of

Oceanography and Fisheries, (Gulfs of Suez and Aqaba Branch) site # 4 and the most northern site # 5, at Suez City. At field, all observations on tidal cycle and changes in animal and plant communities were recorded; while collected specimens were preserved in 10 % seawater formalin solutions.



Figure (1): Shows the locations of the study sites.

At the laboratory, all preserved specimens of porcelain crab were sorted, sexually differentiated and counted, then weighed to the nearest 0.1 gm using an electric balance with accuracy of 0.01 g after removing excess water by blotting with absorbent tissues. Carapace length (CL), carapace breadth (CB), and second male pleopodal length (PL) were measured to nearest mm with a Caliper Verneer with accuracy of ± 0.01 mm. The adult (matures) and small immature or juvenile crabs were dissected, gonads were examined and Chi-square test was used for comparing sex ratios for all population throughout the study period.

The size at first maturity was determined for each sex by applying the relationship between males and females right chela propodal length and male's right second pleopodal length against carapace length according to Hartnoll (1982) and El-Sayed (1997).

The ovigerous (gravid) females were noticed and examined, and all incubated eggs were classified into five developmental stages according to methods of Subramoniam (1982), Schembri (1982) and El-Sayed *et al.*, (1998). The colour and incubation period for these eggs were recorded at each stages of incubation, while the frequencies of ovigerous females were used as an indicator for spawning season according to Subramoniam (1982).

Egg diameters were measured to the nearest micrometer (μm) using Sterio-binocular microscope (PZO Warsaw) provided with graduated screen. Eggs were separated from pleopods, and put in saline solution for two hours, then taken onto a slide and measured under the microscope at power of magnification of 80 X.

Fecundity denotes to the total number of incubated eggs per female. It was determined according to Subramoniam (1982) and Schembri (1982) by counting all incubated eggs carried by ovigerous females after separation from pleopodal setae by a dissecting needle in a divided Petri- dish. The relationship between mean egg number (absolute fecundity, F) of each size class and female carapace length (CL) was calculated according to the following formula:

$$\text{Log F} = \log a \pm b \text{Log CL} \quad (\text{Haynes } et al., 1976)$$

Where F = absolute fecundity, CL = independent variable represented by carapace length in mm, "a" is a constant equal to the intercept of straight line with Y-axis and "b" is an exponent, equal to the slope of fitted line.

Results

A-Sexual dimorphism:

The adults mature individuals of the flattened porcelain crab, *Petrolisthes rufescens* are crab-like, have symmetrical abdomen, folded under thorax. The abdominal appendages are the main sex character for rapid differentiation between mature males and females of this species. However, it was very difficult to distinguish sex in specimens of 3 mm CL or less. Mature males have relatively tapering abdomen provided with two unequal pairs of uniramus pleopods. The second pleopods are well developed and large, compared with very short anterior ones. On contrast, in maturing females abdomen is more flattened, semicircular, provided with four biramus pleopods, usually fringed with entangled setae during spawning season.

On the other hand, chelipeds are slightly unequal in both sexes. Right chela is usually larger than left; appears clearly longer than carapace length in adult males and females, but tend to be slightly larger in males.

B- Sex ratio:

The results in Table (1) show that the overall sex ratio of *P. rufescens* during the present study is 1: 0.70 (males to females) with highly significant Chi-square ($X^2 = 7.486$, $P < 0.05$, $df = 1$). Monthly variations in sex ratio were clearly observed. In spite of the ratios of males to females were 1:1 during June, October and December 2000, the ratios of females were unstable and greatly varied during most months of this study, recording overdominance for males. However, female's ratios increased remarkably to 1:1.17 (males to females) during March 2001, and reached the maximum ratio, was 1:1.40 (males to females) during September 2000, but showing lowest ratio was 1: 0.26 (males to females), recorded in April 2001. The statistical analysis by Chi-square test showed significant differences in sex ratios between different months during this study ($X^2 = 20.185$, $df = 25$, $P < 0.05$).

C- The size at first maturity:

From the relationship between carapace length against right chela propdal length for males and females and right second pleopodal length in males it has been revealed that *P. rufescens* males and females attain sexual maturity between 4.0 and 6.0 mm CL (Figures 2-4). This is very clear from an inflection in the relations for both mature and immature individuals. It means that, all males and females below 4 mm CL are certainly sexually immature, while those have 6 mm CL or above become sexually maturing, but that lie between 4 and 6 mm are mixed from both groups.

Table (1): Total number of collected specimens, percentage of ovigerous females and sex ratios of *P. rufescens* from Ain Sukhna, Gulf of Suez

Months	Number & %	Total No. ♀ ♀ & ♂ ♂	Mature females			% of ovigerous females	Mature ♂ ♂	♀ ♀ & ♂ ♂ Juveniles	Sex ratio (♂ ♂: ♀ ♀)
			All females	Spent	Ovigerous females				
April 2000	47	19	7	12	63.16	28	-	1:0.68	
May	Area did not visit								
June	24	12	6	6	50.0	12	-	1:1	
July	20	9	1	8	88.89	10	1	1:0.90	
August	23	9	3	6	66.67	12	2	1:0.75	
September	27	14	6	8	57.14	10	3	1:1.40	
October	14	7	7	0	0	7	-	1:1	
November	12	3	3	0	0	9	-	1:0.33	
December	11	5	5	0	0	5	1	1:1	
January 2001	14	4	4	0	0	8	2	1:0.50	
February	15	4	2	2	50.0	11	-	1:0.36	
March	13	7	7	0	0	6	-	1:1.17	
April	24	5	2	3	60.0	19	-	1:0.26	
May	12	4	0	4	100	8	-	1:0.50	
Total	256	102	53	49		145	9	1:0.70	




Figure (2): The relationship between male carapace length (CL) and right chela second propodal length (RChPL) of the porcelain crab, *P. rufescens* from Ain Sukhna, Gulf of Suez.




Figure (3): The relationship between female carapace length (CL) and right chela second propodal length (RChPL) of the porcelain crab, *P. rufescens*, from Ain Sukhna, Gulf of Suez.




Fig. 1. Relationship between carapace length and right second pleopodal length of the porcelain crab, *P. rufescens* from Ain Sukhna, Gulf of Suez.

The onset of smallest ovigerous females was also taken for determining the first maturity size. Field collection showed that the smallest ovigerous female measured 4.15 x 4.0 mm (CL x C B), captured in July 2000, which means that, most females attain sexual maturity between 4 and 5 mm CL, and may be become sexually maturing above 5 mm CL. Also, laboratory examination showed that the abdomen of immature females was tightly closed against sternum, with fine, delicate and slender pleopods, fringed with fine and weak setae. It becomes semicircular, provided with strong pleopods, fringed with dense marginal setae in full mature and adult females.

It is also worth to mention that, all individuals of juvenile equal to or below 3 mm CL were not sexually recognizable in their external morphology. It was also difficult to even identify them to their specific taxa due to the presence of juveniles of the coexisting less common porcelain crab species, *Petrolisthes leptocheles* in the same habitats.

D- The incubated eggs and developmental stages:

The incubated eggs or eggs carried with ovigerous females are varied from oval or semi-oval, to elliptical with polygonal surfaces resulting from their crowding on the pleopods. Their overall size was varied according to developmental stages from 400 to 1000 μm with an average of $640 \pm 105 \mu\text{m}$. These eggs were classified into the following five stages:

- Stage I: It denotes to the newly laid eggs after recent ovulation. They appear bright yellow, larger in size, with an average diameter of $810 \pm 70 \mu\text{m}$.
- Stage II: It refers to the early stage of incubated eggs, which have yellow, orange or brown colour, with an average size of $610 \pm 80 \mu\text{m}$.
- Stage III: It denotes to the incubated eggs characterized by the presence of eyed-embryos, of pale orange or gray hues colour, with an average size of $605 \pm 75 \mu\text{m}$,
- Stage IV: This stage refers to incubated eggs having well developed embryos have whitish colour and an average size of $650 \pm 80 \mu\text{m}$,

- Stage V or full complete embryo: It denotes to the releasing larvae, characterized by grey egg-cases attached to pleopod setae, with remaining few eggs with an average size of $640 \pm 70 \mu\text{m}$

It was noticed that, the size of incubated eggs were remarkably decreased gradually with advances in maturation. The average egg size decreased sharply from $810 \pm 70 \mu\text{m}$ for the newly laid eggs, to 610 ± 80 , 605 ± 75 , 650 ± 80 , and $640 \pm 70 \mu\text{m}$ for the other following developmental stages of maturation, respectively, with a slightly increase in stages IV and V.

E- Incubation period:

The present results indicated that, the appearance of ovigerous females carrying eggs at late stages of incubation (complete embryos with eyes and appendages) were recorded from February to September. However, the appearance of all stages varied from the newly laid eggs to late stages with complete embryos during September, and their complete disappearance throughout October, November, December and the following January, implies that these eggs did not exceed than one month for incubation under female's abdomen and determines the incubation period which can be estimated about one month.

F- Spawning season:

The spawning season of this species is determined based on the appearance of berried or ovigerous females, characterizes by carrying clutches of fertilized eggs on their pleopods under her abdomen. The data in Table (1) and Figure (5) show that the porcelain crab, *P. rufescens*, has lengthy definite breeding season, starts often at late winter and early spring and extends through summer to early autumn.

The first appearance for ovigerous females was recorded in February, and its occurrence continued throughout spring and summer to early autumn (end of September). During 2001, the frequencies of ovigerous females increased gradually from 50 % in February, reaching 100 % during May of the same year. The same pattern was recoded in 2000, where ovegrious female's ratios increased from 63.16 % in April, and continued with high frequencies during summer, being 88.89 % in July, but declined to 66.67 % in August, showing more decline to 57.14 % during September, and disappeared completely during the period from October 2000 to the following January 2001 (Table 1 & Figure 5), in spite of an intensive survey which had been done in these months.




Figure (5): Monthly percentages for *Petrolisthes rufescens* ovigerous females showing spawning season.

G - Juveniles recruitments:

Results in Table (1) display the total number of collected specimens including maturing females (spent and ovigerous), maturing males and juveniles of both sexes. In spite of low percentage of juveniles, the data showed that the first appearance of recognizable juvenile of this species was recorded in July 2000 followed by continuous appearance throughout August and September as well as during winter (December 2000 and January 2001). The delaying of juvenile appearance in July in spite recording of late stages of eggs with complete embryos in February denotes to the relative long period for juvenile metamorphosis after hatching.

H -Fecundity:

The incubated eggs for 41 *P. rufescens* ovigerous females were counted. The egg number or fecundity for these females was relatively low, varying from 35 to 450, with an average of 208 ± 92 eggs /female. As previously mentioned, the smallest ovigerous female was 4.15 x 4.0 mm (CL x CB), collected in July 2000, and carrying 48 eggs, all at late stages of maturation with embryos having prominent eyes and well developed appendages. On the other hand, the largest egg number was 450 for female of 8.45 x 8.25 mm (CL x CB) collected in June 2000; whereas the lowest egg number was 35 for female of 4.65 x 4.55 mm (CL x CB), collected in August 2000. It is worth to mention that, 8 ovigerous females were found have egg clutches with very few eggs. These females may be subjected to egg loss. For example, the largest captured ovigerous female attained 9.0 x 8.85 mm (CL x CB), carrying egg clutches of 195 eggs, all were newly laid and at early stage of incubation, but several pleopd setae were ruptured and egg free.

The logarithmic relationship between carapace length (CL) and total egg number or fecundity (F) for 27 ovigerous females (have nearly intact egg clutches) is illustrated in Figure (6), and represented by the following equation:

$$\text{Log F} = 0.455 + 2.21 \text{ Log CL}$$

This relation is statistically significant, and curvilinear with high positive correlation coefficient “r” being 0.82, an allometric regression coefficient “b” 2.21, and positive intercept of Y-axis, “a” = 0.455.




Figure (6): The relationship between carapace length and mean egg number of *P. rufescens* females from Ain Sukhna, Gulf of Suez

Discussion

It is well known that, the reproductive biology of decapod crustaceans is well documented, particularly for brachyurans (Warner, 1977; Sastry, 1983; Lowery and Nelson, 1988; El-Sayed *et al.*, 1998; Fouda, 2000; El-Sayed, 2004). However, little is known about the reproductive biology of Family Porcellanidae (Anomura) around the world (Ahmed and Mustaqim, 1974; Fransozo and Bertini, 2001; Hollebone and Hay, 2007; Monaco *et al.*, 2010) and no information available till now about reproduction of *Petrolithes rufescens* from the entire Red Sea and its associated Gulfs (Suez and Aqaba) in spite of its abundance along the Red Sea coasts (Lewinsohn, 1969; El-Sayed, 2002; Fouda *et al.*, 2003).

The results of the present study showed that the individuals of *Petrolithes rufescens* exhibited morphological sexual dimorphism helps for sex differentiation. The over all sex ratio for this species was 1: 0.7 (males to females), showing overdominance for males on females during most months of the study period, with only an increase in female's ratios during certain months. These results are in well agreement with those mentioned by Oliveira and Masunari (1995) on *Petrolisthes armatus* in Brazil and Hollebone and Hay (2007) on the same species in Gorgia, and that of Farnsozo and Bertini (2001) on *Pachycheles monilefer* belong to family Prcellanidae in São Paulo coast (Brazil), where sex ratios remain nearly close to 1:1 throughout the year, but with seasonal fluctuations. In contrast, the present sex ratios are contradicted with that reported from South Carolina, which were male-biased at a ratio of 1.5:1 (Hartman and Stancyk 2001).

REPRODUCTIVE BIOLOGY OF THE FLATTENED PORCELAIN 27

The sharp fluctuations in female's ratios particularly before or through spawning season may be coincides with the restriction in female's movements in their burrows or escaping into slightly deep water either for egg deposition (ovulation) and hatching, which agrees with results given by Lowery and Nelson (1988), El-Sayed *et al.* (1998), Fouda (2000) and El-Sayed (2004), on true crabs, or due to more catchability of ovigerous females than males during some months of spawning (El-Sayed, 1992, El-Sayed *et al.*, 1998), or attributes to other environmental factors including differential mortality, migration or habitat selection, depth preference, changes in salinity, temperature and other environmental parameters which agrees with Hill (1975), Thomas (1975), and Potter *et al.* (1983) on true crabs and Farnsozo and Bertini (2001), Hollebone (2006), Hollebone and Hay(2007) and Monaco *et al.*, (2010) on porcelain crabs.

Using of relative allometric growth rates for *P. rufescens* individuals before and after puberty moult gave results very similar to that mentioned on other decapods particularly brachyuran species. The present results showed that, all males and females below 4 mm CL are often sexually immature, but those have 6 mm CL or above become sexually maturing. These results are in full agreement with the previous works on porcelanid green crab *Petrolisthes armatus* carried out by Boudreaux *et al.* (2006). The authors showed that collected females were sexually mature at just 3-4 mm carapace width; as well as with those reported by Jones (1977), Scelzo (1985), Oliveira and Masunari (1995), Fransoz and Bertini (2001) and Monaco *et al.* (2010) on species of some porcellaniid genera. These results are also in agreement with that carried out on true crabs by Somerton (1981), Schembri (1982), Potter *et al.*,(1983), Jewett *et al.*,(1985), El-Sayed (1997, 2004), El-Sayed *et al.*, (1998), Fouda (2000).

The progressive increase in male chelae size after maturity is required for the enhancement of defense, feeding and reproductive functions (Hartnoll, 1974; El-Sayed *et al.*, 1998). In the flattened porcelain crab, chelae are used for defense and reproduction functions, but do not employ for feeding (Barnard, 1950; Rypien and Palmer, 2007). On the other hand, broadness of the females abdomen is useful to perform its function for carrying and protecting incubated eggs and helps maintain the continuous circulation of oxygenated water over the eggs carried on her underside (Hartnoll, 1974; Schembri, 1982; Oliveira and Masunari 1995; El-Sayed, 1997, 2004; Hartman and Stancyk, 2001; de Lastang, *et al.*, 2003; Hollebone, 2006),

Like most Red Sea animals, *P. rufescens* population has definite lengthy breeding season, extending usually from February to the following September. It coincides by the appearance of ovigerous females from February to the end of September, and exhibits remarkable seasonality in breeding of this species as known as for most Red Sea invertebrates (El-Sayed, 1992, 1997, 2004; El-Sayed *et al.*, 1998; Fouda, 2000) and agrees well with that mentioned by Ahmed and Mustaquim (1974) on the same species collected from Manora Island, Karachi, Pakistan, where gravid females appear annually from January to November, and with those recorded from subtropical (Fransozo and Bertini, 2001) and temperate decapods (Schembri, 1982).

The use of decapod ovigerous females as tool for determination of breeding seasons in anomurans had been employed in several species such as the tropical hermit crab, *Clibanarius clibanarius* (Varadarajan and Subramoniam, 1982) and hermit crab *Pagurus bernhardus* (Lancaster (1990) as well as in several species of porcelain crabs (Ahmed and Mustaqim, 1974, Fransozo and Bertini, 2001, and Hollebone and Hay, 2007).

The presence of *P. rufescens* ovigerous females with different developmental incubated egg stages throughout the period from February to September may indicate frequencies of spawning during the same breeding season. It also may define the length of incubation period, which may reach to nearly one month, which coincides with the results mentioned by El- Sayed (1997, 2004), El-Sayed *et al.*, (1998), Fouda (2000) on some decapods from the Rd Sea and Gulf of Suez.

Hollebone (2006) explained that when *Petrolisthes armatus* eggs hatch they metamorphose through two planktonic zoeal stages and then one planktonic megalopal stage before finally becoming settlement-ready juveniles. The similar case is observed for *P. rufescens* in the present study where its juveniles are appeared in July, delayed for about 4 months after appearance of ovigerous females in February.

The present results showed that the fecundity of *P. rufescens* is low in comparison with that reported for other intertidal relatively same size decapod species such as *Leptosdium exaratus* from the same area (Fouda, 2000) and those from both the Red Sea (El-Sayed, 2004) and Indian Ocean (Siddiqui & Ahmed, 1992). The decreasing in fecundity of *P. rufescens* from the Gulf of Suez is attributed to several factors including the prevailing hazard environmental conditions, or according to semi-terrestrially habits for this species represented by relative large size of incubated eggs or due to the biological factors dominated by fighting and predations, or restriction of food resources (Warner, 1977).

The presence of good curvilinear relationship between fecundity and carapace length, with an allometric regression coefficient 'b'(2.21), agrees well with that reported by Monaco *et al.*, (2010) on *Petrolisthes granulosus*, Siddiqui and Ahmed (1992), El-Sayed (1997, 2004), El-Sayed *et al.*(1998) and Fouda (2000) on true crabs.

Although the present results provide detailed information about the reproduction of porcelain crab, *P. rufescens*, as one of the most common intertidal decapods along the western coast of the Gulf of Suez, other studies are still required to assess their current status particularly at the same sites after exposing to negative impacts of the expanding constructions of tourist villages, resorts, camp sites and hotels, not only at the Gulf of Suez coasts but also all over the Red Sea Egyptian coasts. Therefore, these results represent the baseline data about this species in Ain Sukhna and can be used as a biological indicator for comparison with that will be taken at the present time after prevailing changes in natural habitats along the Gulf of Suez coasts through the first decade of the twenty first century.

References

1. Ahmed, M. and Mustaqim, J. (1974): Population structure of four species of porcellanid crabs (Decapoda, Anomura) occurring on the coast of Karachi. *Mar. Biol.*, 26: 173- 182.
2. Barnard, K. H. (1950): Descriptive Catalogue of South Africa Decapod Crustacea (Crabs and Shrimps). *Ann. S. Afr. Mus.* 38: 1-400.
3. Boudreaux, M. L.; Stiner, J. L., and Walters, L. J. (2006): Biodiversity of sessile and motile macrofauna on intertidal oyster reefs in Mosquito Lagoon, Florida. *J. Shellfish. Res.*, 25:1079-1089.
4. de Lasting, S.; Hall, N. and Potter, I. C. (2003): Influence of a deep artificial entrance channel on the biological characteristics of the blue swimmer crab *Portunus pelagicus* in a large microtidal estuary. *J. Exp. Mar. Biol. Ecol.* 295(1): 41-61.
5. El-Sayed, A. A. M. (1992): Biological studies on some brachyuran crabs (Crustaceans) from Suez Canal. Ph. D. Thesis. Zool. Dept. Fac. Sci., Al- Azhar University, Cairo.
6. El-Sayed, A. A. M. (1997): The biology of spider crab, *Meneathius monoceros* from the Red Sea and Gulf of Aqaba, Egypt. *Egypt. J. Aquat. Biol. & Fish.*, 1(1):1-16.
7. El-Sayed, A. A. M. (2002): Horizontal and vertical zonation of crustaceans within the black mangrove, *Avicennia marina* (Forskål) Vierh, extending along the Egyptian Red Sea Coast. *Al Azhar Bull. Sci.* 13(2): 121-146.
8. El-Sayed, A. A. M. (2004): Some aspects of the ecology and biology of the intertidal xanthid crab, *Leptodius exaratus* (H. Milne Edwards, 1834) from the Egyptian Red Sea Coast. *J. Egypt. Ger. Soc. Zool.* 45 (D): 115-139.
9. El-Sayed, A. A. M; Saber, S. A., El-Damhougy, K. A. and Fouda, M.M.A. (1998): The reproductive biology of grapsid crab, *Metopograpsus messor* (Forskål,1775) from Ain Sukhna, Gulf of Suez, Egypt. *J. Aquat. Biol. & Fish.* 2(4):359-377.
10. Fouda, M. M. A. (2000): Biological and ecological studies on some crustacean decapods from the Suez Gulf. M. Sci. Thesis, Zoology Department, Faculty of Science, Azhar University, Cairo.
11. Fouda, M. M., El-Sayed, A. A. M. and Fouda, F. M. (2003): Biodiversity and ecology of crustacean fauna in the mangal ecosystems along the North- western Red Sea coast of Egypt. *Al-Azhar Bull. Sci.* 14 (1): 121-142.
12. Fransozo, A. and Bertini, G. (2001): Population structure and breeding period of *Pachycheles monilifer* (Dana) (Anomura, Porcellanidae) inhabiting sabellariid sand reefs from the littoral coast of Sao Paulo, Brazil. *Revta Bras. Zool.*, 18 (1): 197-203.
13. Haig, J. (1960): The Porcellanidae (Crustacea, Anomura) of the eastern Pacific. *Allan Hancock Pacific Expeditions*, 24: 1-440.
14. Hartman, M. J. and Stancyk, S. E. (2001): Distribution of an invasive anomuran decapod, *Petrolisthes armatus*, in the North Inlet-Winyah Bay national Estuarine Research Reserve (NERR) on the South Carolina coast. Abstract. 2nd. *Intern. Conf. on Mar. Bioinva.*, April 9-11, 2001. New Orleans, LA.
15. Hartnoll, R. G. (1974): Variation in growth pattern between some secondary sexual characters in crabs (Decapoda: Brachyura). *Crustaceana* 27(2):131-136.
16. Hartnoll, R. G. (1982): Growth. In: *The Biology of Crustacea*. Vol. II (L. G. Abele, ed.), Academic Press, New York, pp.111-197.

17. Haworth, A. H.(1882): A new binary arrangement of the marurous Crustacea. *The philosophical Magazine and Journal*, 65: 105- 106, 183- 184.
18. Haynes, E., Karinen, J. F.; Watson, J. and Hopson, D. J. (1976): Relation of number of eggs and egg length to carapace width in the brachyuran crabs, *Chionoectes bairdi* and *C. opilio* from the Southeastern Bering Sea amd *C. opilio* from the Gulf of St. Lawrence, south eastern Bering Sea. *J. Fish. Res. Board. Can.* 33:2592-2595.
19. Heller, C. (1961): Beiträge zur Crustaceen- Fauna des rothen Meeres. *Zweiter Theil. Kla. Der kai. Akad. Wiss. Wien* 44(1): 241-295.
20. Hill, B. J. (1975): Abundance, breeding and growth of the crab *Scylla serrata* in two South African Estuaries. *Mar. Biol.* 32:119-126.
21. Hollebhone A. L. (2006): An invasive crab in the South Atlantic Bight: Friend or foe? *Unpublished Doctoral Dissertation. Georgia Institute of Technology.* 133 p.
22. Hollebhone, A. L. and Hay, M. E. (2007): Population dynamics of the non-native crab *Petrolisthes armatus* invading the South Atlantic Bight at densities of thousands m². *Marine Ecology Progress Series* 336: 211- 223.
23. Ismail, I. M. (2005): Ecological and biological studies on some intertidal benthic communities in Suez Gulf, Red Sea, Egypt, M. Sci. Thesis, Zoology Department, Faculty of Science, Al-Azhar University, Cairo.
24. Jewett, S. C.; Solan, N. A. and Somerton, D. A. (1985): Size at sexual maturity and fecundity of the Fjord-dwelling golden king crab, *Lithodes aequispina* Benedict, from northern British Columbia. *J. Crust. Biol.* 5(3): 377-385.
25. Jones, M. B. (1977): Breeding and seasonal population changes of *Petrolisthes elongates* (Crustacea, Decapoda, Anomura) at Kaikoura New-Zeland. *J. Royal Soc. New Zealand*, 7(3): 259- 272.
26. Lancaster, I. (1990): Reproduction and life history strategy of the hermit crab *Pagurus bernhardus*. *J. Mar. Biol. Ass. U. K.* 70: 129- 142.
27. Lewinsohn, Ch. (1969): Die Anomuren Des Roten Meeres (Crustacea: Decapoda: Paguridea, Galatheaidea, Hippidea). *Rijksmuseum van Natuurlijk Historie, Leiden, E. J. Berill, The Netherlands*, 214 pp. + 2 Pls.
28. Lowery, W. A. and Nelson W. G. (1988): Population ecology of the hermit crab *Clibanaris vitatus* (Decapoda: Diogenidae) at Sebastian Inlet Florida USA. *J. Crustacean Biol.*, 8(4): 548-556.
29. Melo, G. A. S. (1999): Manual de identificacáodos Crustacea Decapoda do litoral brasileiro: Anomura, Thalassinidea, Palinuridea e Astacidea. São Paulo, Editora Plêiade, 551 p.
30. McLaughlin, P. A.; Lemaitre, R. and Sorhannus, U. (2007): Hermit crab phylogeny: A reappraisal and its “fall-out”. *J. Crust. Biol.*, 27(1): 97–115.
31. Monaco, C. J.; Brokordt, K. B. and Gaymer, C. F.(2010): Latitudinal thermal gradient effect on the cost of living of the intertidal porcelain crab *Petrolisthes granulatus*. *Aquat. Biol.*, 9: 23-33.
32. Ng, P. K. L.; Guinot, D. and Davie, P. J. F. (2008): Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles Bulletin of Zoology*, Supplement, 17: 1–286.

REPRODUCTIVE BIOLOGY OF THE FLATTENED PORCELAIN 31

33. Oliveira, E. and Masunari, S. (1995): Estrutura populacional de *Petrolisthes armatus* (Gibbes) (Decapoda, Anomura, Porcellanidae) da Ilha do Farol, Matinhos, Paraná, Brasil. *Revta. Bras. Zool.*, 12(2): 355- 371.
34. Osawa, M. and McLaughlin, P. A. (2010): Annoated checklist of anomuran decapod crustaceans of the world (exclusive of the Kiwaoidea and families Chirostylidae and Galatheidae of the Galatheoidea) Part II- Porcellanidae. *The raffles Bulletin of Zoology, Supplement* No. 23: 109- 129.
35. Potter, I. C.; Chrystal, P. J. and Loneragan, N. R.(1983): The biology of the blue crab, *Portunus pelagicus* in an Australian Estuary. *Mar.Biol.* 78:75-85.
36. Rypien, K. L. and Palmer, R. A. (2007): The effect of sex, size and habitat on the incidence of puncture wounds in the claws of the porcelain crab *Petrolisthes cinctipes* (Anomura: Porcellanidae). *J. Crust. Biol.*, 27(1): 59- 64.
37. Sastry, A. (1983): Ecological aspects of reproduction. In: *The Biology of Crustacea*. Vernberg, F. J. & Vernberg, W. (eds.), Vol. 8. *Academic Press, New York*, P. 179- 270.
38. Scelzo, M. A. (1985): Biología y morfometría del canerejo *Petrolisthes politus* (Gray, 1831) (Anomura, Porcellanidae) de la Isla Cubagua, Venezuela. *Biol. Inst. Oceanogr. Univ. Oriente*, 24(1-2): 242- 252.
39. Schembri, P. J. (1982): The biology of a population of *Ebalia tuberosa* (Crustacea: Decapoda: Leucosidae) from the Clyde Sea area. *J. Mar. Biol. Ass.U.K.* 62:101-115.
40. Siddiqui, G. and Ahmed, M. (1992): Fecundities of some marine brachyuran crabs from Karachi (Pakistan). *Pakistan J. Zool.* 24(1):43-45.
41. Somerton, D. A. (1981): Regional variation in the size and maturity of two species of tanner crab *Chionoecetes bairdi* and *C. opelio* in the eastern Bering sea, and its use in defining management subareas. *Can. J. Fish. Aqua. Sci.* 38:163-174.
42. Subramoniam, T. (1982): Determination of reproductive periodicity in the intertidal mole crab, *Emeritta asiatica*. In: *Manual of Research Methods for Marine Invertebrates and Reproduction* (Subramoniam, T. ed.). CMFRI Publication, No.9.Cochin, India, pp.163-167
43. Thomas, M. M. (1975): Reproduction, fecundity, and sex ratio of the green tiger prawn, *Penaeus semisulcatus* de Haan. *Indian J. Fish.*, 21(1): 152-163.
44. Varadarajan, S. and Subramoniam, T. (1982): Reproduction of the continuous breeding tropical hermit crab, *Clibanarius clibanarius*. *Mar. Ecol. Prog. Ser.* 8:197-201.
45. Veloso, V. G. and de Melo, G. A. S (1993): Taxonomía distribuição da família Porcellanidae (Crustacea, Decapoda, Anomura) on litoral brasileiro. *Iheringia, Serie Zoologia*, 75: 171–186.
46. Warner, G. F. (1977): *The Biology of Crustacea*. Elek Science, London.

بيولوجية التكاثر في سرطان البورسيلين، بيتروليسيس روفيسينس (البورسيلينيدي، السرطانات ملتوية البطن، القشريات) من منطقة العين السخنة بخليج السويس، مصر

عواد عبده محمد السيد⁽¹⁾، مصطفى مختار فودة⁽²⁾، أحمد مسعد عزب⁽¹⁾

إسماعيل مسعود إسماعيل⁽³⁾

(1) قسم علم الحيوان، شعبة علوم البحار والأسماك، كلية العلوم بنين، جامعة الأزهر، مدينة نصر، القاهرة، مصر، (2) مستشار بجهاز شئون البيئة، المعادي، القاهرة، مصر، (3) ماجستير علوم البحار.

الملخص العربي

أجريت هذه الدراسة على نوع سرطان البورسيلين *Petrolisthes rufescens* المنتشر في المناطق المدية لشواطئ الساحل الشمالي الغربي لخليج السويس في الفترة من أبريل 2000 إلى مايو 2001.. وقد تم اختيار خمس مناطق لإجراء هذه الدراسة كانت من الجنوب إلى الشمال: (1) جنوب العين السخنة، (2) بلاج الخيمة، (3) بلاج السماد، (4) شاطيء المعهد القومي لعلوم البحار والمصايد بالسويس، (5) شاطيء مدينة السويس. ونظرا لتأثر المناطق الشمالية بكل من الأنشطة البشرية والتلوث البترولي معا كما بالمنطقتين 5&4، أو بزيادة النشاط البشري والمعماري المتمثل بزيادة القرى والمنتجعات السياحة في المنطقتين 2&3 جعل من الصعب متابعة إجراء الدراسات الحقلية بصفة منتظمة بتلك المناطق، لذا فقد تم اختيار المنطقة الأولى (جنوب العين السخنة) لإجراء الدراسات الحقلية والبيئية والبيولوجية بصفة شهرية نظرا لكونها أبعد - حين ذاك عن تلك الأنشطة سالفة الذكر وأقلها نوعا ما أثرا بالنفط.

ولقد أوضحت النتائج العملية والحقلية أن نوع سرطان البورسيلين *P. rufescens* موسم تكاثر ممتد، يرتبط بظهور الإناث حاضنات البيض المخصب يبدأ مبكرا في أواخر الشتاء (فبراير) ويستمر طوال أشهر موسمي الربيع والصيف لينتهي في أوائل الخريف مع نهاية شهر سبتمبر، مع الزيادة الملحوظة في ارتفاع نسب تلك الإناث خاصة خلال فصلي الربيع والصيف، ودخول الأجيال الجديدة إلى بيئة الأطوار اليافعة (الآباء) بدءا من شهر يولييه ويستمر دخولها خلال موسمي الصيف والخريف. كما أشارت النتائج إلى الزيادة الواضحة في نسبة الذكور عن الإناث حيث كانت النسبة الشقية العامة 1: 0.70 (ذكور: إناث)، مع تأرجحها الموسمي واختلاف قيم أحد الشقين عن الآخر باختلاف المواسم. كما بينت النتائج وصول معظم الأطوار غير الناضجة من الذكور والإناث إلى مرحلة النضج الجنسي فيما بين 4- 6 مم من طول الدرقة لتصبح جميعها ناضجة جنسيا في الأطوار الأطول من 6 مم. وتظهر إناث هذا النوع إخصابية منخفضة نوعا ما، إذ تراوح عدد البويضات المحضنة تحت البطن فيما بين 35 و 450 بويضة بمتوسط 208 ± 92 بويضة / أنثى، وأظهرت التحاليل الإحصائية وجود علاقة ارتباط واضحة بين زيادة عدد البويضات وزيادة حجم الإناث. كما أوضحت القياسات أن قطر البويضات المحضنة تراوح فيما بين 400 و 1000 ميكرون بمتوسط عام 640 ± 105 ميكرون، مع ظهور تدرج في اللون مع تقدم مراحل النضج بدء من الأصفر ثم البرتقالي فالبنّي وصولا إلى اللون الرمادي الشاحب لليرقات الكاملة قبيل الانطلاق.

ومما تجدر الإشارة إليه أن منطقة الدراسة بالعين السخنة كانت حينئذ شبه بعيدة عن تأثير الأنشطة البشرية، ولكن شوهد منذ الأشهر الأخيرة من الدراسة امتداد تلك الأنشطة إليها متمثلة في ردم الكثير من الأماكن الصخرية من أجل زيادة مواقع الاصطياف وكذلك تأثرها بزيادة التلوث النفطي مما يجعل هذه الدراسة بمثابة قاعدة بيانات يمكن الرجوع إليها لتقييم الوضع الراهن عن تلك المنطقة خاصة بعد أن أصبحت الآن تعج بالكثير من القرى السياحية وتعرضها للتلوث النفطي.