

6-1-2015

Section: Mathematics, Statistics, Computer Science, Physics and Astronomy

Evaluation of the Elliptic Integrals

Sharaf M.A.

1Department of Astronomy, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

Alrawjih F.A.

Department of Mathematics, Faculty of Science, University of Dammam, Dammam Saudi Arabia

Follow this and additional works at: <https://absb.researchcommons.org/journal>



Part of the [Life Sciences Commons](#)

How to Cite This Article

M.A., Sharaf and F.A., Alrawjih (2015) "Evaluation of the Elliptic Integrals," *Al-Azhar Bulletin of Science*: Vol. 26: Iss. 1, Article 10.

DOI: <https://doi.org/10.21608/absb.2015.22632>

This Original Article is brought to you for free and open access by Al-Azhar Bulletin of Science. It has been accepted for inclusion in Al-Azhar Bulletin of Science by an authorized editor of Al-Azhar Bulletin of Science. For more information, please contact kh_Mekheimer@azhar.edu.eg.

Evaluation of the Elliptic Integrals

M.A. Sharaf¹ and F.A. Alrawjih²

¹Department of Astronomy, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

²Department of Mathematics, Faculty of Science, University of Dammam, Dammam Saudi Arabia

Abstract:

The present paper is devoted for establishing accurate computational algorithms for the incomplete and complete elliptic integrals (EI) of the first, second and third kind. For these goals, we first derived some properties of EI that could be used to check the validity and the accuracy of the algorithms; in addition, particular continued fraction expansion of the ratio of the complete elliptic integrals of the second and first kinds is also derived. Secondly, we established the trigonometric series expansions of EI, together with the recurrence formulae of their coefficients so as to facilitate the computations. Also, Gautschi's algorithm of the top-down continued fraction evaluation is described. Numerical applications are performed for: (a) the incomplete elliptic integrals using their trigonometric series expansions, (b) the complete elliptic integrals of the second kind from the complete elliptic integrals of the first kind using Gautschi's algorithm. Finally the numerical results were checked by two ways:

- i- by satisfying the conditions given by properties of EI.
- ii- by comparing their values with those list in slandered tables.

In this respect, the numerical results show excellent arguments with these ways, a fact which proves the validity, accuracy and the effeteness of our algorithms

Keywords:

Elliptic integrals, recursive computations algorithms, continued fraction, trigonometric series expansions.

1. Introduction

It can be shown that, if $R(x,y)$ is a rational function of x and y and $g(x)$ is a polynomial in x of degree 3 or 4, then the integral

$$\int R\{x, \sqrt{g(x)}\} dx,$$

can be expressed as a linear combination of terms, each of which is either an elementary function, or an elliptic integral of the first, second, or third kind. Accordingly, numerical evaluation of a few integrals permits precise evaluation of a broad class of integrals. On the other hand the elementary problem in analytical mechanics – the motion of the simple pendulum – cannot be accurately described without resorting to elliptic integrals. Moreover, elliptic integrals play an important role in many aspect of Astrodynamics, e.g. the gravitational attraction of a solid homogeneous ellipsoid upon an exterior particle is represented in terms of elliptic

integrals. Again the period of rectilinear oscillation [1] of an infinitesimal mass along the straight line through the center of mass and perpendicular to the plane of rotation of the finite masses is a linear combination of complete elliptic integrals of the first, second, and third kinds. Also the planetary orbit determination in Einstein's gravitational theory, leads to an elliptic integral [2]. In fact there are many applications of the elliptic integral e.g. [3].

Due to their importance as mentioned briefly in the above, the present paper is devoted for establishing accurate computational algorithms for the incomplete and complete elliptic integrals (EI) of the first, second and third

2. Basic Formulations

2.1 Definitions

The normal forms of the elliptic integrals F and E of the first and second kind, respecti

$$\text{are } F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} = \int_0^{\sin \phi} \frac{dx}{\sqrt{(1-x^2)(1-x^2 k^2)}}, \quad (1)$$

$$E(k, \phi) = \int_0^\phi \sqrt{1-k^2 \sin^2 \phi} \, d\phi = \int_0^{\sin \phi} \sqrt{\frac{1-x^2 k^2}{1-x^2}} dx, \quad (2)$$

which are functions of the amplitude ϕ , where $0 < \phi \leq \pi/2$ and of the modulus k , where $0 \leq k \leq 1$, (The symbol m , called the parameter, is also used in place of the square of the modulus, i.e. $m \equiv k^2$). The first form of these special functions is called Legendre's form. The alternate,

$$F(k, \pi/2) \equiv K(k) \equiv K \quad \text{and} \quad E(k, \pi/2) \equiv E(k) \equiv E. \quad (3)$$

Otherwise, they are referred to as incomplete elliptic integrals.

The elliptic integral of the third kind is

$$\Pi(\nu, k, \phi) = \int_0^\phi \frac{d\phi}{(1-\nu \sin^2 \phi) \sqrt{1-k^2 \sin^2 \phi}} = \int_0^{\sin \phi} \frac{dx}{(1-\nu x^2) \sqrt{(1-x^2)(1-x^2 k^2)}}, \quad (4)$$

where ν is the characteristic. The properties of the integral depend on the location of the characteristic.

obtained from the first by setting $x = \sin \phi$ is called Jacobi's form.

When the amplitude $\phi = \pi/2$, the integrals are then complete elliptic integrals and denoted by $K(k)$ and $E(k)$ or simply K and E . Thus

2.2 Some basic properties of the hypergeometric function

In the subsequent analysis we need some basic properties of the hypergeometric function $y = F_1(\alpha, \beta, \gamma; x)$ which is defined as

$$F_1(\alpha, \beta, \gamma; x) = \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k} \frac{x^k}{k!}, \quad (5)$$

where

$$(\eta)_0 = 1; \quad (\eta)_k = \eta (\eta+1) (\eta+2) \dots (\eta+k-1). \quad (6)$$

Among the properties of $y = F_1(\alpha, \beta, \gamma; x)$, we need the following basic three properties:

1-Gauss's differential equation for the hypergeometric function $y = F_1(\alpha, \beta, \gamma; x)$ is:

$$x(1-x) \frac{d^2 y}{dx^2} + [\gamma - (\alpha + \beta + 1)x] \frac{dy}{dx} - \alpha \beta y = 0. \quad (7)$$

2- Gauss's identity of the hypergeometric function $y = F_1(\alpha, \beta, \gamma; x)$ is:

$$\gamma(1-x) F_1(\alpha, \beta, \gamma; x) = \gamma F_1(\alpha-1, \beta, \gamma; x) - (\gamma-\beta) x F_1(\alpha, \beta, \gamma+1; x). \quad (8)$$

3- The ratio $F_1(\alpha, \beta+1, \gamma+1; x) / F_1(\alpha, \beta, \gamma; x)$ is given as a continued fraction of the form

$$\frac{F_1(\alpha, \beta+1, \gamma+1; x)}{F_1(\alpha, \beta, \gamma; x)} = \frac{1}{1 - \frac{\delta_1 x}{1 - \frac{\delta_2 x}{1 - \frac{\delta_3 x}{1 - \dots}}}}, \quad (9)$$

where

$$\delta_{2n+1} = \frac{(\alpha+n)(\gamma-\beta+n)}{(\gamma+2n)(\gamma+2n+1)}, \quad \delta_{2n} = \frac{(\beta+n)(\gamma-\alpha+n)}{(\gamma+2n-1)(\gamma+2n)}. \quad (10)$$

2.3 Some properties of the elliptic integrals

In what follows some properties of the elliptic integrals will be listed, while their proofs are given in Appendix A. These properties could be used to check the

validity and the accuracy of the computational algorithms which used for the valuations of IE, as

1- For fixed amplitude, we have

$$\phi \leq F(k, \phi) \leq \ln(\tan \phi + \sec \phi), \quad (11)$$

$$\sin \phi \leq E(k, \phi) \leq \phi. \quad (12)$$

2- $F(k, \phi)$ and $E(k, \phi)$ are both odd functions of ϕ , i.e.

$$F(k, -\phi) = -F(k, \phi), \quad (13)$$

$$E(k, -\phi) = -E(k, \phi). \quad (14)$$

3- $F(k, \phi)$ and $E(k, \phi)$ satisfy the identities

$$F(k, n\pi + \phi) = 2nK(k) + F(k, \phi), \quad (15)$$

$$E(k, n\pi + \phi) = 2nE(k) + E(k, \phi), \quad (16)$$

where n is any positive integer.

4- The special case of the elliptic integral of the third kind for which the characteristic ν is equal to the parameter $m = k^2$ is given by

$$(1 - k^2) \Pi(k^2, k, \phi) = E(k, \phi) - \frac{k^2 \sin 2\phi}{2\sqrt{1 - k^2} \sin^2 \phi}. \quad (17)$$

5- The complete elliptic integrals $K(m)$ and $E(m)$ of the first and second kind, satisfy the following differential equations

$$2m(1 - m) \frac{dK}{dm} = E - (1 - m)K, \quad (18)$$

$$2m \frac{dE}{dm} = E - K, \quad (19)$$

$$m(1 - m) \frac{d^2K}{dm^2} + (1 - 2m) \frac{dK}{dm} - \frac{1}{4}K = 0, \quad (20)$$

$$m(1 - m) \frac{d^2E}{dm^2} + (1 - m) \frac{dE}{dm} + \frac{1}{4}E = 0. \quad (21)$$

6- The complete elliptic integrals $K(m)$ and $E(m)$ could be expressed in terms of the hypergeometric functions F_1 as

$$K(m) = \frac{\pi}{2} F_1\left(\frac{1}{2}, \frac{1}{2}, 1; m\right), \quad (22)$$

$$E(m) = \frac{\pi}{2} F_1\left(-\frac{1}{2}, \frac{1}{2}, 1; m\right), \quad (23)$$

where $m \equiv k^2$.

7- The complete elliptic integrals $K(m)$ and $E(m)$ satisfy the identity

$$(1 - m) \frac{K(m)}{E(m)} = 1 - \frac{m}{2} \frac{F_1\left(\frac{1}{2}, \frac{1}{2}, 2; m\right)}{F_1\left(\frac{1}{2}, -\frac{1}{2}, 1; m\right)}, \quad (24)$$

8- The complete elliptic integral $E(m)$ of the second kind could be expressed in terms of the complete elliptic integral $K(m)$ of the first kind by the continued fraction expansion as

$$E(m) = \frac{(1 - m)K(m)}{1 - \frac{m}{2 - \frac{3m}{4 - \frac{m}{2 - \frac{5m}{8 - \frac{3m}{2 - \frac{7m}{12 - \frac{5m}{2 - \dots}}}}}}}}}. \quad (25)$$

3. Trigonometric Series Expansions of the Incomplete Elliptic Integrals

In this section, the trigonometric series expansions of the elliptic integrals of the first second and third kind will be established.

3.1 The incomplete elliptic integral of the first kind

Since

$$F(k, \phi) = \int_0^{\phi} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi, \quad (26)$$

then by using the binomial theorem we get

$$F(k, \phi) = \sum_{n=0}^{\infty} (-1)^n \binom{-\frac{1}{2}}{n} k^{2n} s_n, \quad (27)$$

where $s_n = \int_0^{\phi} \sin^{2n} \phi d\phi$. Expressing $\sin^{2n} \phi$ as a series of cosine multiples of ϕ , we get

$$s_n = 2^{-2n} \left\{ \binom{2n}{n} \phi + \sum_{m=1}^n (-1)^m \binom{2n}{n-m} \frac{\sin 2m\phi}{m} \right\}. \quad (28)$$

3.2 The incomplete elliptic integral of the second kind

By the same way as in the above section we get

$$E(k, \phi) = \sum_{n=0}^{\infty} (-1)^n \binom{\frac{1}{2}}{n} k^{2n} s_n, \quad (29)$$

where s_n is given by Equation (28).

3.3 The incomplete elliptic integral of the third kind

Recalling the definition of the elliptic integral of the third kind as

$$\Pi(\nu, k, \phi) = \int_0^{\phi} (1 - \nu \sin^2 \phi)^{-1} (1 - k^2 \sin^2 \phi)^{-1/2} d\phi. \quad (30)$$

Expand each term of the integrand using the binomial theorem, perform the product of the resulting power series of $\sin^{2q} \phi$, then expressing $\sin^{2q} \phi$ as a series of cosine multiples of ϕ , we get

$$\Pi(\nu, k, \phi) = \sum_{n=0}^{\infty} \nu^n s_n \sum_{m=0}^n (-1)^m \binom{-\frac{1}{2}}{m} \left(\frac{k^2}{\nu} \right)^m. \quad (31)$$

4. Computational Developments

The computational developments for the EI will be considered through the following points.

4.1 Computation forms of the equations

For the computational purposes the above infinite series should be truncated to $(N+1)$ terms. So, we have

$$F(k, \phi) = \sum_{n=0}^N T_n s_n, \quad (32)$$

$$E(k, \phi) = \sum_{n=0}^N H_n s_n, \quad (33)$$

$$\Pi(\nu, k, \phi) = \sum_{n=0}^N G_n s_n, \quad (34)$$

where

$$T_n = (-1)^n \binom{-\frac{1}{2}}{n}, \quad (35)$$

$$H_n = (-1)^n \binom{\frac{1}{2}}{n}, \quad (36)$$

$$G_n = \nu^n \sum_{m=0}^n T_m \left(\frac{k^2}{\nu} \right)^m. \quad (37)$$

4.2 Recurrence formulae

In what follows, some recurrence formulae for the coefficients will be established to facilitate their computations.

From equations (35) and (36) we get for the T 's and H 's coefficients the recurrence formulae

$$T_{n+1} = \frac{1}{2}(2n+1)T_n; \quad T_0 = 1, \quad (38)$$

$$H_{n+1} = \frac{1}{2}(2n+1)H_n; \quad H_0 = 1. \quad (39)$$

From Equation (37) we have

$$\begin{aligned} G_{n+1} &= \nu^{n+1} \sum_{m=0}^{n+1} T_m \left(\frac{k^2}{\nu} \right)^m \\ &= \nu^{n+1} \left\{ T_{n+1} \left(\frac{k^2}{\nu} \right)^{n+1} + \sum_{m=0}^n T_m \left(\frac{k^2}{\nu} \right)^m \right\} \\ &= \nu^{n+1} T_{n+1} \left(\frac{k^2}{\nu} \right)^{n+1} + \nu G_n, \end{aligned}$$

consequently

$$\frac{G_{n+1} - \nu G_n}{G_n - \nu G_{n-1}} = \nu^n \left(\frac{T_{n+1}}{T_n} \right) \left(\frac{k^2}{\nu} \right) \xrightarrow{\text{Using Equ.(38)}} = \frac{1}{2}(2n+1)k^2,$$

from which we get the recurrence formula

$$G_{n+1} = \frac{1}{2}G_n \left\{ 2\nu + (2n+1)k^2 \right\} - \frac{1}{2}k^2 \nu (2n+1)G_{n-1}, \quad (40)$$

while its initial values are obtained from Equations (35) and (37) and we get

$$G_0 = 1; \quad G_1 = \nu + \frac{1}{2}k^2. \quad (41)$$

Finally, a recurrence formula for the s function could be obtained as follows. Write s_n as

$$s_n = \int_0^{\phi} \sin^{2n-1} \phi \sin \phi d\phi,$$

integrating by parts we deduce that

$$(2n-1)s_{n-1} - 2ns_n = \cos \phi \sin^{2n-1} \phi,$$

also

$$(2n+1)s_n - 2(n+1)s_{n+1} = \cos \phi \sin^{2n+1} \phi.$$

From these two equations we get the recurrence formula

$$s_{n+1} = \frac{1}{2(n+1)} \left\{ [2n + 2n \sin^2 \phi + 1] s_n - (2n-1) \sin^2 \phi s_{n-1} \right\}, \quad (42)$$

while its initial values are

$$s_0 = \phi; \quad s_1 = \frac{1}{2} \phi - \frac{1}{4} \sin 2\phi. \quad (43)$$

4.3 Continued fraction method

In fact, continued fraction expansions are generally far more efficient tools for evaluating the classical functions than the more familiar infinite power series. Their convergence is typically faster and more extensive than the series.

4.3.1 Top-Down Continued Fraction Evaluation

There are several methods available for the evaluation of continued fraction. Traditionally, the fraction was either computed from the bottom up, or the numerator and denominator of the n^{th} convergent were accumulated separately with three-term recurrence formulae. The

draw back of the first method is obviously, having to decide far down the fraction to being in order to ensure convergence. The draw back to the second method is that the numerator and denominator rapidly overflow numerically even though their ratio tends to a well defined limit. Thus, it is clear that an algorithm that works from top-down while avoiding numerical difficulties would be ideal from a programming standpoint.

Gautschi [4] proposed very concise algorithm to evaluate continued fraction from the top-down and may be summarized as follows. If the continued fraction is written as

$$c = \frac{n_1}{d_1 + \frac{n_2}{d_2 + \frac{n_3}{d_3 + \ddots}}},$$

then initialize the following parameters

$$a_1 = 1, \quad b_1 = n_1/d_1, \quad c_1 = n_1/d_1;$$

and iterate ($k = 1, 2, \dots$) according to

$$a_{k+1} = \frac{1}{1 + \left[\frac{n_{k+1}}{d_k d_{k+1}} \right] a_k}, \quad b_{k+1} = [a_{k+1} - 1] b_k, \quad c_{k+1} = c_k + b_{k+1}.$$

In the limit, the c sequence converges to the value of the continued fraction.

Continued fraction method was used in many problems in astrophysics [e.g. 5 & 6] as well as in the special functions of Astrodynamics [e.g. 7 & 8].

4.4 Numerical applications

1. Consider the 30 random values of ν, ϕ (radian) and m ($\equiv k^2$) listed in Table (1) of Appendix B. Numerical applications of the analytical formulations of subsections 4.1 and 4.2 yield for the incomplete elliptic integrals EI1, EI2 and EI3 of the first, second and third kind respectively the values listed in Table (2) of Appendix B.
2. The complete elliptic $E(m)$ ($m \equiv k^2$) of the second kind could be computed in terms of the complete elliptic integral $K(m)$ of the first kind by the continued fraction expansion of Equation (25). For this purpose we used Gautschi's algorithm with $n_1 = (1 - m) K, \quad n_2 = -m, \quad n_i$ (i odd) $= -i m, \quad n_i$ (i even) $= -(i - 3) m,$
 $d_1 = 1, \quad d_j$ (j odd) $= 2j - 2, \quad$ and d_j (j even) $= 2,$

$$\text{Min}[F(k, \phi)] = \int_0^\phi d\phi = \phi,$$

$$\text{Max}[F(k, \phi)] = \int_0^\phi \frac{d\phi}{\sqrt{1 - \sin^2 \phi}} = \int_0^\phi \sec \phi \, d\phi = \ln(\tan \phi + \sec \phi);$$

$$\text{Max}[E(k, \phi)] = \int_0^\phi d\phi = \phi,$$

$$\text{Min}[E(k, \phi)] = \int_0^\phi \sqrt{1 - \sin^2 \phi} \, d\phi = \sin \phi;$$

which prove the inequalities (11), (12).

Formulae (13) & (14):

Equations (13) and (14) follow directly from the definitions of $F(k, \phi)$ and $E(k, \phi)$.

Formulae (15) & (16):

We can write Equations (1) and (2) as

$$F(k, \phi = \int_0^{\pi/2} (1 - k^2 \sin^2 \psi)^{-1/2} \, d\psi + \int_{\pi/2}^\pi (1 - k^2 \sin^2 \psi)^{-1/2} \, d\psi - \int_\phi^\pi (1 - k^2 \sin^2 \psi)^{-1/2} \, d\psi,$$

$$E(k, \phi = \int_0^{\pi/2} (1 - k^2 \sin^2 \psi)^{1/2} \, d\psi + \int_{\pi/2}^\pi (1 - k^2 \sin^2 \psi)^{1/2} \, d\psi - \int_\phi^\pi (1 - k^2 \sin^2 \psi)^{1/2} \, d\psi.$$

In the second and third integrals, we let $\theta = \pi - \psi$, then $\phi = -\phi$, so we get by using Equations (13) and (14), that

$$F(k, \pi + \phi) = 2 K(k) + F(k, \phi),$$

$$E(k, \pi + \phi) = 2 E(k) + E(k, \phi).$$

These are Equations (15) and (16) which are true for $n = 1$.

In the above equations we let $\phi = \phi + \pi$, we get

$$F(k, 2\pi + \phi) = 2 K(k) + F(k, \pi + \phi) = 4 K(k) + F(k, \phi),$$

$$E(k, 2\pi + \phi) = 2 E(k) + E(k, \pi + \phi) = 4 E(k) + E(k, \phi).$$

These are Equations (15) and (16) which are true for $n = 2$.

so, we get the values of $E(m)$ for some m as listed in Table (3) of Appendix B.

Finally the numerical results were checked by two ways:

- i- by satisfying the conditions given by properties of EI.
- ii- by comparing their values with those list in slandered tables.

In this respect, the numerical results show excellent arguments with these ways, a fact which proves the validity, accuracy and the effeteness of our algorithms.

Appendix A

The Proofs of the Formulae of Subsection 2.3

Formulae (11) & (12):

Since $F(k, \phi)$ and $E(k, \phi)$ are continuous functions, then

$$\text{Min}[F(k, \phi)] \leq F(k, \phi) \leq \text{Max}[F(k, \phi)],$$

and

$$\text{Min}[E(k, \phi)] \leq E(k, \phi) \leq \text{Max}[E(k, \phi)].$$

From Equations (1) & (2) we have

Assume Equations (15) and (16) which are true for n , so

$$F(k, n\pi + \phi) = 2n K(k) + F(k, \phi),$$

$$E(k, n\pi + \phi) = 2n E(k) + E(k, \phi);$$

then let $\phi = \phi + \pi$, in these two equations we get

$$F\{k, (n+1)\pi + \phi\} = 2n K(k) + F(k, \pi + \phi) = 2(n+1) K(k) + F(k, \phi),$$

$$E\{k, (n+1)\pi + \phi\} = 2n E(k) + E(k, \pi + \phi) = 2(n+1) E(k) + E(k, \phi).$$

These are Equations (15) and (16) which are true for $n = n + 1$, since they are true for $n = 1, n = 2, \dots$, consequently they are for any n positive integer.

Formula (17):

$$\text{Let } y = \sqrt{1 - k^2 \sin^2 \phi},$$

$$\text{then } y' = \frac{-k^2 \sin 2\phi}{2\sqrt{1 - k^2 \sin^2 \phi}},$$

$$\text{and } \frac{dy^2}{dx^2} = y'' = -T_1 - T_2, \quad (\text{A-1})$$

$$\text{where } T_1 = \frac{k^2 \cos 2\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \frac{k^2}{\sqrt{1 - k^2 \sin^2 \phi}} + 2\sqrt{1 - k^2 \sin^2 \phi} - \frac{2}{\sqrt{1 - k^2 \sin^2 \phi}},$$

$$T_2 = \frac{k^4 \sin^2 2\phi}{4(1 - k^2 \sin^2 \phi)^{3/2}} = \frac{-k^2 \cos^2 \phi}{(1 - k^2 \sin^2 \phi)^{1/2}} + \frac{k^2 \cos^2 \phi}{(1 - k^2 \sin^2 \phi)^{3/2}},$$

$$\text{that is } T_2 = \frac{-k^2}{(1 - k^2 \sin^2 \phi)^{1/2}} - (1 - k^2 \sin^2 \phi)^{1/2} + \frac{2}{(1 - k^2 \sin^2 \phi)^{1/2}} + \frac{k^2 - 1}{(1 - k^2 \sin^2 \phi)^{3/2}}.$$

Using the above expressions for T_1 and T_2 into Equation (A-1), and then integrating the resulting equation, so we get

$$(1 - k^2) \Pi(k^2, k, \phi) - E(k, \phi) = \frac{k^2 \sin 2\phi}{2\sqrt{1 - k^2 \sin^2 \phi}},$$

which is Equation (17).

Formulae (18):

$$\text{Since } K(m) = \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - m \sin^2 \phi}},$$

differentiating with respect to m we get

$$\frac{dK}{dm} = -\frac{1}{2m} K(m) + \frac{1}{2m} \Pi(m, m, \pi/2),$$

then using Equation (17) with $\phi = \pi/2$, we get Equation (18).

$$\text{Formulae (19) : Since } E(m) = \int_0^{\pi/2} \sqrt{1 - m \sin^2 \phi} d\phi,$$

differentiating with respect to m we get

$$\frac{dE}{dm} = \frac{1}{2m} \int_0^{\pi/2} \frac{1 - m \sin^2 \phi - 1}{\sqrt{1 - m \sin^2 \phi}} d\phi = \frac{1}{2m} (E - K),$$

then is Equation (19).

Formulae (20):

Differentiating Equation (18) and then using Equation (19) for dE/dm and Equation (18) for $(K - E)/2m$ we get Equation (20).

Formulae (21):

Differentiating Equation (19) and then using Equation (18) for dK/dm and Equation (19) for K , we get Equation (21).

Formulae (22):

Comparing Equation (20) with Gauss's Equation (7) for hypergeometric function we get, when

$$y \equiv K, x \equiv m, \gamma = 1 \Rightarrow \alpha + \beta = 1, \alpha \beta = 1/4 \Rightarrow \alpha = 1/2, \beta = 1/2,$$

consequently
$$K(m) = C_1 F_1\left(\frac{1}{2}, \frac{1}{2}, 1; m\right).$$

Satisfies Gauss's equation. Since $K(0) = \pi/2$ and $F_1\left(\frac{1}{2}, \frac{1}{2}, 1; 0\right) = 1$, then $C_1 = \pi/2$ and Equation (22) is proved.

Formulae (23):

Comparing Equation (21) with Gauss's Equation (7) for hypergeometric function we get when $y \equiv E, x \equiv m, \gamma = 1 \Rightarrow \alpha + \beta = 0, \alpha\beta = -1/4 \Rightarrow \alpha = -1/2, \beta = 1/2$,

consequently

$$K(m) = C_2 F_1\left(-\frac{1}{2}, \frac{1}{2}, 1; m\right).$$

That satisfies Gauss's equation. Since $E(0) = \pi/2$ and $F_1\left(-\frac{1}{2}, \frac{1}{2}, 1; 0\right) = 1$, then $C_2 = \pi/2$ and Equation (23) is proved.

Formulae (24):

Let $\gamma = 1, \alpha = 1/2, \beta = 1/2$ and $x \equiv m$

in Equation (8), and then divide both sides of the resulting equation by $F_1\left(-\frac{1}{2}, \frac{1}{2}, 1; m\right)$ and using Equations (22)

& (23), then Equation (24) follows immediately.

Formulae (25):

Let $\gamma = 1, \alpha = 1/2, \beta = -1/2$ and $x \equiv m$

in Equation (9) an using Equation (10) for the δ 's we get

$$Q = \frac{F_1\left(\frac{1}{2}, \frac{1}{2}, 2; m\right)}{F_1\left(\frac{1}{2}, -\frac{1}{2}, 1; m\right)} = \frac{1}{1 - \frac{3m/8}{1 - \frac{m/8}{1 - \frac{5m/16}{1 - \dots}}}}$$

Using Equation (24) for $E(m)$, Equation (25) yields directly.

Appendix B
Numerical Results

Table (1): Some values of ν, ϕ (radian) and m ($\neq k^2$).

No.	ν	ϕ	m	No.	ν	ϕ	m
1	0.886722	0.866325	0.594413	16	0.371839	1.077740	0.617374
2	0.619108	0.918401	0.427062	17	0.152632	0.203947	0.264464
3	0.355766	1.019590	0.539380	18	0.196058	0.852554	0.619489
4	0.402134	0.978971	0.218435	19	0.453673	0.955426	0.805967
5	0.104672	0.422304	0.501638	20	0.867869	0.386516	0.316444
6	0.722051	1.058850	0.310770	21	0.235311	0.883901	0.431224
7	0.648157	0.318158	0.682142	22	0.321819	0.291290	0.892084
8	0.712991	0.414826	0.518002	23	0.661250	0.757539	0.299885
9	0.375125	0.361961	0.527260	24	0.810326	0.550609	0.625692
10	0.141408	0.685071	0.388569	25	0.248589	0.117576	0.736812
11	0.422924	0.245789	0.615998	26	0.602711	0.572889	0.665022
12	0.797856	0.282745	0.739104	27	0.405484	0.937949	0.660505
13	0.610132	0.975660	0.682898	28	0.508192	0.771637	0.607257
14	0.510288	0.729252	0.206240	29	0.243918	0.895272	0.435174
15	0.143886	1.159740	0.817152	30	0.780660	0.714040	0.554252

Table (2): The values of the three incomplete integrals for the input values of Table (1).

No.	EI1	EI2	EI3	No.	EI1	EI2	EI3
1	0.933299	0.807440	1.224670	16	1.214580	0.965883	1.401850
2	0.972148	0.869631	1.174260	17	0.204313	0.203569	0.204744
3	1.116500	0.936493	1.260500	18	0.919637	0.793721	0.962597
4	1.009400	0.950075	1.143340	19	1.091690	1.847205	1.265270
5	0.428625	0.416148	0.431257	20	0.389534	0.383540	0.407324
6	1.114580	1.007690	1.520750	21	0.932513	0.839556	0.988356
7	0.321859	0.314533	0.329100	22	0.295029	0.287636	0.297739
8	0.421020	0.408793	0.439145	23	0.778307	0.737722	0.890671
9	0.366147	0.357859	0.372217	24	0.568427	0.533769	0.620661
10	0.705534	0.665635	0.720589	25	0.117776	0.117377	0.117911
11	0.247321	0.244275	0.249444	26	0.594392	0.552746	0.636896
12	0.285559	0.279981	0.291818	27	1.035650	0.855554	1.170130
13	1.091190	0.880298	1.356850	28	0.819890	0.728231	1.911368
14	0.741775	0.717093	0.813214	29	0.946269	0.848894	1.006710
15	1.426770	0.971483	1.514440	30	0.748335	0.682450	0.867226

Table (3): $E(m)$ for some m as computed from Gautschi's algorithm.

m	$E(m)$	m	$E(m)$
0.807939	1.17311	0.168876	1.50222
0.863506	1.13337	0.226277	1.47776
0.651545	1.26983	0.539360	1.33055
0.458234	1.37138	0.408748	1.39525
0.314211	1.43898	0.714159	1.23317
0.565508	1.31689	0.581473	1.30841
0.314999	1.43862	0.888529	1.11403
0.274959	1.45650	0.691199	1.24689
0.939370	1.10154	0.910660	1.09592
0.983505	1.02436	0.773534	1.19600
0.707370	1.23726	0.125024	1.52048
0.419361	1.39019	0.139654	1.51443
0.958540	1.05199	0.552426	1.32376
0.861890	1.13458	0.559323	1.32014
0.938420	1.07145	0.559516	1.32004

References

1-Battin, R.H.: 1999, "An Introduction to the Mathematics and Methods of Astro-dynamics, 2nd Edition", AIAA Education Series. USA.

2-Lanczos, C.: 1986, "The Variational Principles of Mechanics" 4th Edition, Dover Publications, Inc. New York.

3-Cayley A.: "An Elementary treatise on Elliptic Functions" 2nd Edition, Dover Publications, Inc. New York.

4-Gautschi, W.: 1967, "Computational Aspects of Three-term Recurrence Relations", SIAM Review, Vol. 9, No. 1, January.

5-Sharaf, M.A.: 2006, "Computations of the Cosmic Distance Equation" Appl. Math. Comput. 174, 1269-1278.

6-Sharaf, M.A.; Almleaky, Y. M; Malawi, A.A; Goharji, A.A and Basurah, H.M.: 2004, "Symbolic Analytical Expressions of the Physical Characteristics for N-Dimensional Radially Symmetrical Isothermal Models", AJSE, King Fahd, Univ., Vol. 29. No. 1A, 67-82.

7-Sharaf, M.A. and Banajh, M.A.: 2001, "Continued Fraction Evaluation of the Stumpff Functions of Space Dynamics", Sci. J. Fac., Minufiya Univ., Vol. XV, 267-282.

8-Sharaf, M.A. and Najmuldeen, S.A.: 2001, "Continued Fraction Evaluation of the Normal Distribution Function", Sci. J. Fac., Minufiya Univ., Vol. XV, 311-324.

9-Abramowitz, M and Stegun I.A. (Editors): 1972, "Handbook of Mathematical Functions", Dover Publications, Inc. New York