

12-1-2016

Section: Botany, Microbiology and Zoology

PLANT COMMUNITIES OF AL TA'AKR MOUNTAIN AND EL SOHOL AREAS, IBB GOVERNORATE, REPUBLIC OF YEMEN

MAREI HAMED

BOTANY AND MICROBIOLOGY DEPARTMENT, AL AZHAR UNIVERSITY, FACULTY OF SCIENCE

Follow this and additional works at: <https://absb.researchcommons.org/journal>



Part of the [Life Sciences Commons](#)

How to Cite This Article

HAMED, MAREI (2016) "PLANT COMMUNITIES OF AL TA'AKR MOUNTAIN AND EL SOHOL AREAS, IBB GOVERNORATE, REPUBLIC OF YEMEN," *Al-Azhar Bulletin of Science*: Vol. 27: Iss. 2, Article 2.

DOI: <https://doi.org/10.21608/absb.2016.24349>

This Original Article is brought to you for free and open access by Al-Azhar Bulletin of Science. It has been accepted for inclusion in Al-Azhar Bulletin of Science by an authorized editor of Al-Azhar Bulletin of Science. For more information, please contact kh_Mekheimer@azhar.edu.eg.

PLANT COMMUNITIES OF AL TA'AKR MOUNTAIN AND EL SOHOL AREAS, IBB GOVERNORATE, REPUBLIC OF YEMEN

MAREI, A . HAMED

AL AZHAR UNIVERSITY, FACULTY OF SCIENCE, BOTANY AND MICROBIOLOGY DEPARTMENT

ABSTRACT

El Sohol area is lie north of Ibb city and Al Ta'akr mountain of Ibb with altitude 3000 m above sea level and lie south of Ibb city. This study is a phytosociological comparison between two areas which differ from each other in topography: A total of 111 species belonging to 51 families of were recorded in 29 sites of the studied area. Family Asteraceae was the most abundant family, comprising 12 species. The families Acanthaceae and Euphorbiaceae represented by 6 species, each Lamiaceae represented by 5 species and Mimosaceae. Moraceae and Poaceae were represented by 4 species.

The plant life form of the studied area showed that the highest recorded life form was for Phanerophytes constituted species representing 37 % of the total species followed by the chaemophytes with 28 species representing 25 %. Therophytes with 26 species representing 23 % . Cryptophytes with 13 species representing 11 % and Hemicryptophytes with only 2 species representing 1.8 % of total species of the work . From AlTa'akr mountain we can summarize the following: Community in plateau dominated by *Pennisetum setaceum*, near plateau was dominated by *Rosa abyssinica* in medium and high slopes, also very steep slopes were dominated by *Hypericum revolutum*, Wadi El Ganat community *Arundo donax* was dominated by in wadi bed, *Ziziphus spina-christi* dominated in the terraces , *Acacia melifera* and *Acacia etbaica* dominated in slopes. In Najed El El Bard, *Ziziphus spina-christi* dominated in wadi bed , *Acacia asak* dominated in the terraces of wadi, *Acacia etbaica* dominated in low slopes of wadi, and *Acacia melifera* dominated in medium slope of wadi. In El Gabal El Akhader . *Ormocarpum dhofarensis* dominated the low slopes *Euphorbia inarticulate* dominated in medium slope and plateau of mountain, while *Cynodon dactylon* high slopes. DECORANA was used to clarify relations among environmental factors and plant distribution. ANOVA was used to detect the most significant factors affecting species distribution. Moisture soil content and calcium content were the most significant factors in determining different ecological groups with P value = 0.000, followed by chlorides with P value = 0.003 and organic carbon with P value = 0.014 .

INTRODUCTION

It is to be mentioned that Ibb governorate is the greenest region of the Republic of Yemen and is known for its heavy rainfall, water springs and basins. It is also famous for growing coffee, incense trees, sugar cane and many important crops. Beside the most famous Wadies in Ibb e.g (Wadi Bana and Wadi Al-Door) there are also high mountains e.g (Sumarah and AlTa'akr) which makes it such an interesting research point. Ibb governorate has an area of 5253 km² with 20 districts. Aqlan (2008) . Several authors write on the flora of Ibb and described species according to elevation but they not give any information about vegetation cover and communities (Forsskål, 1775). (Lavranos, 1971a, b, c). (Wood, 1997). (Miller and Cope, 1996). Al-Khulaidi & Pole (1989). And Al-Hubaishi and Mueller (1984) . Aqlan, (2008) surveyed the flora of Ibb governorate by visiting different localities representing various topographical features of the governorate and collected 416 species belong to 294 genera

and 95 plant families . Ibb governorate lacks plant communities studies . Recently, Marei , et al. (2016) studied plant communities of Badan mountain of Ibb governorate using modern programs for analysis and described the plant communities in Badan mountain.

The present the work is comparative study of the plant communities of AlTa'akr mountain and El Sohol area of Ibb governorate according to their ecology and vegetation analysis and inter-related edaphic factors.


THE STUDY AREA

Ibb governorate is called the green province and it belongs to four topographical units of the Republic of Yemen. These units are western mountains, medium altitude of western mountains, high altitude mountains and highland plains. Ibb governorate consists of twenty districts. These districts extend between (long. 43 o . (Map 1) the study area of the present work including El Sohol area which lie at north from Ibb city and AlTa'akr mountain (3000 m above

sea level.) at south of Ibb city (Map 2). Climatic data of Ibb governorate were obtained as (Statistical year book, 2001-2004). Temperature of Ibb averages between 27.9 °C in January and 33 °C in June which represents a limited variability and a moderate temperature along the year. High vegetation density of the area is due to the high precipitation rate. No precipitation was recorded in January, while from October to February precipitation have been recorded. The maximum precipitation rate was recorded in June with 287.5 mm. Records of wind speed showed the highest speed in December with 5.2 knot/s and the lowest in January with 3.4 knot/s. The highest relative humidity records were in August with 69.75%, while the lowest was in March with 47.25%.



Map 1. Administrative map of Yemen showing different governorates of Yemen Republic.



Map 2. Showing El Sohol area (north Ibb city) and Al Ta'aker mountain (south Ibb city).

MATERIALS AND METHODS

Twenty nine sites were selected to represent the vegetation of AlTa'akrmountain and El Sohol area, sites from 1 to8 were located from 2400 to 3100 m above sea level. Elevation of AlTa'akrmountain, El Sohol area (elevation from 1800 to 2100m above sea level) included the following A: Wadi El Ganat slopes , terraces and wadi bed sites from 9 to 14 , B: Najed El Bard sites from 15 to 24 and C: El Gabal El Akhdar sites from 25 to 29. These sites were subjectively chosen at locations where either dense vegetation or change in species composition was encountered. For each stand, three soil samples were collected at a depth of 5-30 cm. These samples were then pooled together to form one composite sample. Part of the soil sample was used to determine the soil moisture content after Piper (1950). The rest of the soil sample was air-dried, sieved through 2mm mesh and stored at room temperature for further analyses. Electrical conductivity (EC) and pH were determined in the filtrate of 1:5 soil / distilled water (w/v) extract. Organic carbon (O.C.) was measured according to the method of Piper (1950). Carbonates and chlorides were determined using the method described by Jackson (1967). Calcium and magnesium were determined according to the procedure of Richards (1954). Vegetation of ten quadrates (10×10 m) in each stand was investigated. Floristic composition of the 29 sites were recorded. All plant species existing in each stand were listed after complete identification according to Täckholm (1974), Wood (1997) and Boulos (1999-2009). Voucher herbarium specimens were prepared and kept in the herbarium of the Department of Biology, Faculty of Science, Ibb University. According to Shukla and Chandel (1989) Density (D), percentage of frequency (F), abundance (A), cover (C), relative density (RD), relative frequency (RF), relative abundance (RA), relative cover (RC) and importance value (IV) were calculated for each species in each stand. Data management was performed using PC-ORD ver. 5 (McCune and Mefford, 1999) to run TWINSPLAN and DECORANA. (ter Braak, 1988). Analysis of variance (ANOVA) using SPSS ver. 22 (Santoso, 2014) was used to test the most significant soil factors affecting the classification of samples.

RESULTS

A total of 111 species belonging to 51 families were recorded in 29 site of the studied area Family Asteraceae was the most abundant family comprising 12species. The families of Acanthaceae and Euphorbiaceae were represented by 6 species Lamiaceae and Mimosaaceae representing by 5 species. Moraceae and Poaceae was represent by 4species. The plant

life form of the studied area showed that the highest life form recorded was Phanerophytes comprising 37 % of the total species followed by the Chaemophytes with 28 species (representing 25 %of total species). Therophytes with 26 species representing 23%, Cryptophytes with 13 species representing 11 % and Hemicyptophytes with only 2 species representing 1.8 % of total species of the work (Table 1).

Table 1:Plant species recorded in Ibb (Al Ta'akr mountain and El Sohol) P = Present. m= meter elevation above sea level.

Family	Species	Life form	Ibb		Abbreviation in TWINSPAN
			Al Ta'akr mountain 3000 m	AlSohol 1900 m	
Acanthaceae	<i>Barleria trispinosa</i> (Forssk.) Vahl.	Phanerophyte.		P	<i>Bartri</i>
	<i>Blepharis ciliaris</i> (L.) B.L. Burt.	Hemicyptophyte.		P	<i>Blecil</i>
	<i>Hypoestes forskalii</i> (Vahl.) R. Br.	Chamaephyte.		P	<i>Hypfor</i>
	<i>Justicia flava</i> (Vahl.) Vahl.	Chamaephyte.		P	<i>Jusfla</i>
	<i>Ruellia patula</i> Jacq.	Hemicyptophyte	P	P	<i>Ruepat</i>
	<i>Acanthus arboreus</i> Forssk.	Phanerophyte.	P	P	<i>Acaarb</i>
Actinopteridaceae	<i>Adiantum capillus-veneris</i> L.	Cryptophyte	P		<i>Adicap</i>
Agavaceae	<i>Agave sisalana</i> Perr.	Phanerophyte.		P	<i>Agasis</i>
Aloeaceae	<i>Aloe inermis</i> Forssk.	Cryptophyte		P	<i>Aloine</i>
.Amaranthaceae	<i>Aerva javanica</i> (Burm. F.) Juss. ex Schult.	Chamaephyte.		P	<i>Aerjav</i>
	<i>Amaranthus lividus</i> L <i>Alternanthera pungens</i> Kunth.	Therophyte. Therophyte.	P	P P	<i>Amaliv</i>
Apiaceae	<i>Ferula communis</i> L.	Chamaephyte.	P		<i>Fer com</i>
Apocynaceae	<i>Adenium obesum</i> (Forssk.) Roem. & Schult.	Phanerophyte	P P	P	<i>Adeobe</i>
Asclepiadaceae	<i>Caralluma cicatricosa</i> (Deflers) N. E. Br.	Chamaephyte.		P	<i>Carcic</i>
	<i>Ceropegia rupicola</i> Defl.	Chamaephyte.		P	<i>Cerrup</i>
	<i>Kanahia laniflora</i> (Forssk.) R. Br.	Phanerophyte.		P	<i>Kanlan</i>
Asparagaceae	<i>Asparagus falcatus</i> L.	Phanerophyte.	P	P	<i>Aspfal</i>
Asphodelaceae	<i>Kniphofia sumarae</i> Deflers	Cryptophyte.	P		<i>Kni sum</i>
Asteraceace	<i>Echinops spinosissimus</i> Turra	Chamaephyte.	P		<i>Echspi</i>
	<i>Kleinia anteuphorbium</i> (L.) DC	Chamaephyte.		P	<i>Kleant</i>
	<i>Osteospermum vaillantii</i> (Decne.) T. Norlindh	Therophyte	P	P	<i>Ost vai</i>
	<i>Senecio hadiensis</i>	Phanerophyte.	P	P	<i>Senhad</i>

	<i>Tagetes minuta</i> L.	Therophyte		P	<i>Tagmin</i>
	<i>Xanthium strumarium</i> L.	Therophyte	P	P	<i>Xanstr</i>
	<i>Sonchus oleraceus</i> L.	Therophyte	P		<i>Sonole</i>
	<i>Pluchea dioscoridis</i> (L.) DC.	Phanerophyte.		P	<i>Pludio</i>
	<i>Centaurea pseudosinaica</i> Czerep.	Chamaephyte.		P	<i>Cenpse</i>
	<i>Centaurothamnus maximus</i> (Forssk.) Wagenitz & Dittrich	Phanerophyte.	P		<i>Cenmax</i>
	<i>Psiadia punctulata</i> (DC.) Vatk <i>Flaveria trinervia</i> (Spreng.) Mohr.	Phanerophyte. Therophyte	P P	P	<i>Psipun</i> <i>Fla tri</i>
Boraginaceae	<i>Heliotropium europaeum</i> L.	Therophyte		P	<i>Heleur</i>
	<i>Trichodesma ehrenbergii</i> Schweinf. ex Boiss.	Therophyte	P		<i>Triehr</i>
Burseraceae	<i>Commiphora habessinica</i> (Berg.) Engl.	Phanerophyte.		P	<i>Com hab</i>
Cactaceae	<i>Opuntia ficus-indica</i> (L.) Miller	Phanerophyte.		P	<i>Opu fic</i>
Caeslpinaceae	<i>Ceratonia siliqua</i> L.	Phanerophyte.		P	<i>Cersil</i>
	<i>Senna italica</i> Miller	Chamaephyte.		P	<i>Sen ita</i>
	<i>Senna occidentalis</i> (L.) Link	Therophyte			<i>Sen occ</i>
Campanulaceae	<i>Campanula edulis</i> Forssk.	Chamaephyte.		P	<i>Cam edu</i>
Capparaceae	<i>Cadaba farinose</i> Forssk.	Phanerophyte.		P	<i>Cad far</i>
Chenopodiaceae	<i>Chenopodium murale</i> L.	Therophyte.		P	<i>Che mur</i>
Convolvulaceae	<i>Convolvulus arvensis</i> L.	Chamaephyte.		P	<i>Conarv</i>
	<i>Seddera arabica</i> (Forssk.) Choisy.	Phanerophyte	P		<i>Sed ara</i>
	<i>Ipomoea purpurea</i> (L.) Roth.	Therophyte		P	<i>Ipo pur</i>
Caryophyllaceae	<i>Spergularia rubra</i> (L.) J&C. Prosl.	Therophyte	P		<i>Spe rub</i>
Cyperaceae	<i>Cyperus rotundus</i> L.	Cryptophye		P	<i>Cyprot</i>
Dipsacaceae	<i>Scabiosa columbaria</i> L.	Chamaephyte.	P		<i>Scacol</i>
Dracaenaceae	<i>Sansevieria ehrenbergii</i> Schweinf. ex Baker.	Cryptophyte.	P	P	<i>San ehr</i>

Euphorbiaceae	<i>Euphorbia ammak</i> Schweinf.	Phanerophyte		P	<i>Eup amm</i>
	<i>Euphorbia cactus</i> Ehrenb. ex Boiss.	Phanerophyte		P	<i>Eupcac</i>
	<i>Euphorbia hirta</i> L.	Therophyte.	P		<i>Euphir</i>
	<i>Euphorbia peplus</i> L.	Therophyte		P	<i>Euppep</i>
	<i>Euphorbia inarticulate</i> Schweinf	Phanerophyte		P	<i>Eupina</i>
	<i>Jatropha curcas</i> L.	Phanerophyte		P	<i>Jatcur</i>
Fabaceae	<i>Indigofera spinosa</i> Forssk.	Chanerophyte.		P	<i>Indspi</i>
	<i>Indigofera hochstetteri</i> Bak.	Therophyte.		P	<i>Indhoc</i>
	<i>Ormocarpum dhofarensis</i> Hillcoat & Gillett	Phanerophyte	P	P	<i>Ormdho</i>
	<i>Cadia purpurea</i> (Picc.) Ait.	Phanerophyte		P	<i>Cadpur</i>
Geraniaceae	<i>Geranium arabicum</i> Forssk.	Chamaephyte.	P	P	<i>Gerara</i>
	<i>Geranium biuncinatum</i> Kokwaro	Therophyte.	P		<i>Gerbiu</i>
Hypericaceae	<i>Hypericum revolutum</i> Vahl.	Phanerophyte	P		<i>Hyp rev</i>
Lamiaceae	<i>Thymus laevigatus</i> Vahl.	Chamaephyte.	P	P	<i>Thy lae</i>
	<i>Plectranthus barbatus</i> Andr.	Chamaephyte.	P	P	<i>Plebar</i>
	<i>Mentha longifolia</i> (L.) L.	Chamaephyte.	P	P	<i>Menlon</i>
	<i>Mentha Piperia</i> L.	Chamaephyte.	P		<i>Men pip</i>
	<i>Morrubium vulgare</i> L.	Chamaephyte.	P		<i>Mor vul</i>
Malvaceae	<i>Abutilon fruticosum</i> Guill. & Perr.	Phanerophyte.		P	<i>Abu fru</i>
	<i>Hibiscus deflersii</i> Schweinf. ex Cufod.	Phanerophyte		P	<i>Hibdef</i>
	<i>Hibiscus trionum</i> L.	Therophyte	P		<i>Hibri</i>
	<i>Malva parviflora</i> L.	Therophyte.		P	<i>Malpar</i>
Mimosaceae	<i>Acacia asak</i> (Forssk.) Willd.	Phanerophyte Phanerophyte		P	<i>Aca asa</i>
	<i>Acacia etbaica</i> Schweinf.	Phanerophyte		P	<i>Acaetb</i>
	<i>Acacia mellifera</i> (Vahl.) Benth	Phanerophyte		P	<i>Acamel</i>
	<i>Acacia origena</i> Asfaw	Phanerophyte		P	<i>Aca ori</i>
Moraceae	<i>Dorstenia foetida</i> (Forssk.) Schweinf.	Chamaephyte.		P	<i>Dor foe</i>
	<i>Ficus populifolia</i> Vahl.	Phanerophyte		P	<i>Ficfoe</i>
	<i>Ficus cordata</i> Thunb.	Phanerophyte		P	<i>Fic car</i>
Nyctaginaceae	<i>Mirabilis jalapa</i> L.	Chamaephyte	P	P	<i>Mirjal</i>
	<i>Commicarpus</i>	Chamaephyte.		P	<i>Com gra</i>

	<i>grandiflorus</i> (A. Rich.) Standl.				
Oleaceae	<i>Jasminum grandiflorum</i> L.	Phanerophyte.		P	<i>Jas gra</i>
Orchidaceae	<i>Holothrix arachnoidea</i> (A. Rich.) Reichb. f.	Cryptophyte.	P		<i>Hol ara</i>
Orobanchaceae	<i>Cistanche phelypaea</i> (L.) Cout.	Cryptophyte.		P	<i>Cisphe</i>
	<i>Orobanche ramose</i> L.	Cryptophyte.	P		<i>Oro ram</i>
Oxalidaceae	<i>Oxalis corniculata</i> L.	Chamaephyte.	P	P	<i>Oxacor</i>
Papaveraceae	<i>Argemone mexicana</i> L.	Chamaephyte.		P	<i>Arg mex</i>
Poaceae	<i>Arundo donax</i> L.	Cryptophyte		P	<i>Aru don</i>
	<i>Cynodon dactylon</i> (L.) Pers	Cryptophyte	P	P	<i>Cyndac</i>
	<i>Pennisetum setaceum</i> (Forssk .) Chiov .	Cryptophyte	P	P	<i>Penvil</i>
	<i>Hopochne vulgare</i> L .	Cryptophyte	P		<i>Hop vul</i>
Polygonaceae	<i>Emex spinosa</i> (L.) Campd.	Therophyte.	P		<i>Eme spi</i>
	<i>Rumex nervosus</i> Vahl.	Phanerophyte	P	P	<i>Rum ner</i>
	<i>Persicaria salicifolia</i> (Willd.) S. F. Gray	Chamaephyte.	P		<i>Per sal</i>
Portulacaceae	<i>Portulaca oleracea</i> L.	Therophyte.		P	<i>Porole</i>
Primulaceae	<i>Anagallis arvensis</i> L.	Therophyte.	P		<i>Ana arv</i>
Resedaceae	<i>Caylusea hexagyna</i> (Forssk.) M. L. Green	Chamaephyte.	P		<i>Cayhex</i>
	<i>Ochradenus baccata</i> Del.	Phanerophyte		P	<i>Ochbac</i>
Rhamnaceae	<i>Ziziphus spina-christi</i> (L.) Willd.	Phanerophyte		P	<i>Zizspi</i>
Rosaceae	<i>Rosa abyssinica</i> Lindley	Phanerophyte	P		<i>Ros aby</i>
Sapindaceae	<i>Dodonaea viscosa</i> (L.) Jacq.	Phanerophyte		P	<i>Dod vis</i>
Scrophulariaceae	<i>Kickxia elatine</i> (L.) Dum.	Therophyte.	P		<i>Kicela</i>
Selaginellaceae	<i>Selaginella yemensis</i> (Swartz) Spring	Cryptophyte	P		<i>Sel yem</i>
Solanaceae	<i>Solanum glabratum</i> Dunal.	Phanerophyte	P	P	<i>Solgla</i>
	<i>Solanum incanum</i> L. var. <i>incanum</i>	Phanerophyte	P	P	<i>Soline</i>
	<i>Solanum nigrum</i> L.	Therophyte.	P	P	<i>Solnig</i>
	<i>Withania somnifera</i> (L.) Duna	Chamaephyte.		P	<i>Wit som</i>
	<i>Datura innoxia</i> Mill	Therophyte		P	<i>Dat inn</i>
Verbenaceae	<i>Lantana camara</i> L.	Phanerophyte	P	P	<i>Lancam</i>
Vitaceae	<i>Cissus quadrangularis</i> L.	Phanerophyte.		P	<i>Cisqua</i>
	<i>Cissus rotundifolia</i> (Forssk.) Vahl.	Phanerophyte.		P	<i>Cisrot</i>
Zygophyllaceae	<i>Tribulus terrestris</i> L.	Therophyte.		P	<i>Tri ter</i>

The analysis of the vegetation was carried out using 29 sites distributed in the two studied areas as the follows Al-Ta'akr mountain (sites from 1 to 8) from the top to bottom of the slope , El Sohol area included the following : A - Wadi ElGanat slopes , terraces and wadi bed sites from 9 to 14 , B- Najed ElBard sites from 15 to 24 and C- ElGabal ElAkhdar sites from 25 to 29.

AlTa'akr Mountain

From Al-Ta'akr mountain data can summarize the following:

Community of the plateau was dominated by *Pennisetum setaceum* , Communities of the near plateau was dominated by *Rosa abyssinica*. Communities of medium and high slopes also steep slopes were dominated by *Hypericum revolutum*.

El Sohol area(sites from 9 to 29) included the following:

A - Wadi El Ganat

From wadi El Ganat survey the following could be found

Communities dominated in wadi bed were dominated by *Arundo donax*. Communities dominated in the terraces were dominated by *Ziziphus spina-christi*. Communities of slopes of wadi were dominated by *Acacia melifera* and *Acacia etbaica*

B- Najed El Bard

From Najed El El Bard data one can summarize the following:

Communities of the wadi bed were dominated by *Ziziphus spina-christi*. Communities of the terraces of wadi were dominated by *Acacia asak*. Communities of the low slopes of wadi were dominated by *Acacia etbaica*.

Communities of moderate slope of wadi were dominated by *Acacia melifera*. Communities of high and steep slopes were dominated by *Euphorbia cactus* and *Commiphora habessinica*., while communities of plateau of mountain were dominated by *Ormocarpum dhofarensense* and *Kleinia odora* .

El Gabal El Akhader

From El Gabal El Akhader data one summarize the following:

Communities present in low slopes were dominated by *Ormocarpum dhofarensense*. Communities at moderate slopes and plateau of mountain were dominated with *Euphorbia inarticulate*. Communities in high slopes were dominated by *Cynodon dactylon*. And communities of plateau of mountain dominated by *Ormocarpum dhofarensense* and *Kleinia odora*(Table 2 and plates 1,2 and 3).

Table (2) Communities types of the study area (Al Ta'akr mountain sites from 1 to 8 , Wadi El Ganat site from 9 to 14 , Najed El Bard sites from 15 to 24 and El Gabal El Akhader sites from 25 to 29) .

Site No.	Habitat	Dominant	Co Dominant
1	Moderate slope	<i>Rosa abyssinica</i>	<i>Acanthus arboreus</i>
2	Moderate steep slope	<i>Rosa abyssinica</i>	<i>Acanthus arboreus</i>
3	high steep slope	<i>Hypericum revolutum</i>	<i>Rosa abyssinica</i>
4	High danger slope	<i>Hypericum revolutum</i>	<i>Pennisetum villosum</i>
5	Very high slope	<i>Hypericum revolutum</i>	<i>Pennisetum villosum</i>
6	Very high slope	<i>Hypericum revolutum</i>	<i>Rosa abyssinica</i>
7	Near plateau of mountain	<i>Rosa abyssinica</i>	<i>Emex spinosa</i>
8	Plateau of mountain	<i>Pennisetum villosum</i>	<i>Rosa abyssinica</i>
9	Wadi terrace	<i>Ziziphus spina-christi</i>	<i>Acacia melifera</i>
10	Terrace and low slope of mountain	<i>Acacia melifera</i>	<i>Hibiscus deflersii</i>
11	Moderate slope	<i>Acacia etbaica</i>	<i>Cissus rotundifolia</i>
12	Steep slope and terrace	<i>Acacia etbaica</i>	<i>Pennisetum villosum</i>
13	Wadi bed	<i>Arundo donax</i>	<i>Ceratonia siliqua</i>

14	slope	<i>Acacia melifera</i>	<i>Acanthus arboreus</i>
15	Terrace	<i>Acacia asak</i>	<i>Commiphora habessinica</i>
16	High slope	<i>Acacia etbaica</i>	<i>Kleinia odora</i>
17	Plateau of mountain	<i>Ormocarpum dhofarens</i>	<i>Kleinia odora</i>
18	Very danger high slope	<i>Euphorbia cactus</i>	<i>Acacia asak</i>
19	High slope	<i>Commiphora habessinica</i>	<i>Senna etalica</i>
20	High slope	<i>Commiphora habessinica</i>	<i>Acacia etbaica</i>
21	Wadi bed	<i>Ziziphus spina-christi</i>	<i>Jatropha curcas</i>
22	Low slope	<i>Acacia origana</i>	<i>Euphorbia inarticulate</i>
23	Moderate slope of mountain	<i>Acacia melifera</i>	<i>Euphorbia inarticulate</i>
24	Plateau of mountain	<i>Kleinia odora</i>	<i>Acacia melifera</i>
25	Plateau of mountain	<i>Euphorbia inarticulate</i>	<i>Cadia purpurea</i>
26	High slope	<i>Cynodon dactylon</i>	<i>Euphorbia inarticulate</i>
27	Moderate slope	<i>Euphorbia inarticulate</i>	<i>Commiphora habessinica</i>
28	Moderate slope	<i>Euphorbia inarticulate</i>	<i>Acacia melifera</i>
29	mild slope	<i>Ormocarpum dhofarens</i>	<i>Commiphora habessinica</i>



Plate 1. Shows *Hypericum revolutum* community aslo *Rosa abyssinica* appear here as a co dominate species in the high slope of Al Ta'akr mountain.

Plate 2. Shows *Jatropha curcas* and *Argemone mexicana* in Wadi terrace of Najed El Bard.





Plate 3.Shows Al Gabal El Akhader with high slope covered with *Cynodon dactylon*.

Out of the TWINSpan classification, the 29 stands were classified into four groups after three levels of classification. Negative group including sites from 9 - 28 this is large group which classified in second classification into two group negative group (group A) including sites of wadi El Ganat sites 16 , 17 19, 20 22, 24 some exception here this group contain site 21 of Najed El Bard and Negative group (14 sites) which also classified in third level into Negative group

(Group B) including 7 sites 16 . 17 , 19 , 20 24 and site 29 of El Gabal Al Akhader Positive group (Group C) 7 sites this is mixed group which containing some sites of Najed El Bard (15 , 18 and 23) as well as some sites of El Gabal Al Akhadr (25 , 26 , 27 and 28). Positive group (Group D) with eight stands (1 to 8) at the mountain of AlTa'akr where *Rosa abyssinica* . *Hypericum revolutum* and *Pennisetum setaceum* are the dominant species table 3.

Table 3. TWINSpan classification of 29 sites showing different groups of the study area

		11111211122221122222	
		90123416790249583567812345678	
12	Mir jal	1----2-----1-----	000
19	Oxa cor	-----2-----1-----	000
23	Xan str	2--2--3-----1-----	001
27	Sol inc	12-3----4-----1--1---1-----	001
35	Ziz spi	745-539-----1-----	001
36	Aca mel	69---95-----7--89-5-7-----	001
37	Aga sis	1--1-2-----	001
38	Fla tri	2---2-2-2-----	001
40	Cer sil	3--262-----	001
41	Abu fru	2-----	001
42	Arg mex	3-----	001
43	Wit som	1-----	001
44	Hyp for	2-----	001
45	Alt pun	1-----	001
47	Hib def	-7-3---2--2--2-----	001
48	Bel cil	-955-3---2-2-----	001
49	Cis qua	--4-----2-----2-----	001
50	Cis rot	--73--2-----2-2---3-----	001
51	Aer jav	--2-----1-----	001
52	Hib vit	--5--2-----1---	001
53	Aca etb	--98---9--9-----6-----	001
54	Ade obe	---3---12-----	001
55	Jat cur	3--55-8-----	001
56	Alo ine	---2-----1-----	001
57	Opu fic	---2-3-----43-----	001

MAREI, A. HAMED

58	Lan cam	-----2-----1-----	001
59	Bar tri	-----5-----3-----4-----	001
60	Dod vis	-----2-----	001
61	Fic sal	-----1-----	001
62	Fic pop	-----2-----	001
63	Kan lan	-----2-----2-----	001
65	Dat ino	-----1-----	001
80	Comerc	-----4-----4-----	001
90	Com gra	-----2-----2-----	001
98	Ama liv	-3-----	001
99	Bar bi	---2-----	001
101	Aru don	-----9-----	001
102	Cad far	-----5-----	001
103	Ind spi	-----2-----	001
110	Men pip	-----2-----	001
11	Eup hir	-----7-----1-----	010
13	Rue pat	-----2--5-4343-33533-41-1-----	010
39	Sen ita	22---22--55-2-23-----	010
64	Eup ina	-----1-----223--7-89-----	010
66	Aca asa	-----5-----7--3-----	010
67	Com hab	-----9-6962-2263-----	010
68	Kle odo	-----8843-89334-2-----	010
69	Ind spi	-----33-3--22---2-----	010
70	Eup cac	-----4-----38--7-4-----	010
71	Tag mim	-----2-----1-3-----	010
72	Tri ter	-----2-----	010
73	Car cic	-----21--2-31-----	010
74	Orm dho	-----59-7-49-4-315-1---2---	010
75	Jus fla	-----364343--3543-----	010
76	Cyp rot	-----2---2-----	010
77	Kle ant	-----6-----	010
78	Sed ara	-----9-----2	010
79	Com for	-----2-----1-----	010
81	Eup amm	-----9-----	010
82	Cad pur	-----75-8-----	010
83	Che mur	--2-----3--2-----	010
84	Jas gra	-----223-----	010
85	Och bac	-----2-----	010
100	Aca ori	----3---4--9-----	010
104	Ind hoc	-----2--23-----	010
106	Hel eur	-----1-----	010
111	Cer rup	-----2-----	010
18	Sol gla	-----22--3--1-----1---12-	011
93	Ple bar	1-----3-----3321-3---22-2	011
95	Men lon	-2-----3--2--2-2-----4-	011
105	San ehr	-----3--2---3-2-----4-	011
20	Cyn dac	3-----567--74---9	100
1	Ros aby	-----98555668	101
2	Aca arb	-----2-----483--2-	101
3	Hyp qua	-----359999--	101
5	Thy lae	-----2--2456-252	101
6	Res sph	-----221-4--2	101
7	Eme spi	-----2-1--23-	101
8	Ech spi	-----1-1--221	101
9	Tri aff	-----1---2--	101
10	Ana arv	-----1-1-----	101
14	Cen max	-----1-1---3-	101
21	Psi pun	-----2-----2--3--31--	101
22	Sca col	-----3-2232	101
26	Cen pse	-----1-----	101
28	Hol arz	-----22---	101
29	Tec gem	-----22---	101

30	Spo rub	-----2---	101
31	Fer com	-----1131	101
32	Kic sca	-----12-	101
34	Fun sp	-----2-----2324---	101
86	Adi cap	-----1-----	101
87	Kni sum	-----2-1343--	101
88	Cay hfx	-----2-----	101
89	Pie bar	-----2-----	101
91	Sel yem	-----32---	101
92	Per sal	-----2-----	101
94	Ger biu	-----22---	101
96	Ger ara	-----2-----2-	101
107	DOr foe	-----1-----	101
108	Hop sch	-----2-1-----	101
109	Mor vul	-----1-----	101
4	Rum ner	---32-----3-2---	110
15	Sol neg	1-----1-----	110
16	Asp fal	--2---2-----21-2--2--	110
17	Pen vil	---62-----3--57--	110
24	Ost vai	1-----1-----	110
25	Plu dis	22-----1-----1--2-2	110
46	Ama gra	1--1-----3--	110
97	Ipo pur	---3-----2-	110
33	Sen had	1-----1-----1--	111

000000000000000000000000111111111

0000000111111111111111111111111

0000000111111111

***** TWINSpan completed *****

After TWINSpan classification, data was analyzed by DECORANA (indirect gradient analysis). Results showed that sites of El Sohol area (the plateau) were positively correlated to axis 2 represented by groups A, B and C, while group D was related to axes 1 including sites from 1 to 8 were located in El Taaker mountain peak were positively correlated to axes 1. Ordination results were in accordance with TWINSpan classification which is quite common in vegetation studies. Axis 1 group D was dominated by *Rosa abyssinica*. *Hypericum revolutum* and *Pennisetum setaceum* on the other hand, groups correlated to axis 2 Group A was dominated by *Ziziphus spina-christi*, *Acacia*

etibca, *Acacia milefera*, and *Arundo donxia*, Group B was dominated by *Acacia etibca*. *Acacia origana*, *Ormocarpum dhofarensis*, *Commiphora habessinica*, *Euphorbia inarticulate* and *Kleinia odora*, and Group C was dominated by *Acacia asak*, *Euphorbia cactus*, *Cordia purpurea*, *Acacia melifera*, *Cynodon dactylon*, and *Euphorbia inarticulate*. Group D was related to axes 1 including sites from 1 to 8 which representing Al Ta'akr mountain peak were positively correlated to axes 1. Soil parameters correlation to axis 1 and 2 were also examined. Moisture content, Na⁺, Ca⁺⁺, and organic carbon were positively correlated to axis 1, while pH, Cl⁻, Co³⁻, K⁺, Mg⁺⁺ were positively correlated to axis 2. (Fig1).




Fig. 1. DCA ordination diagram for the vegetation groups of the 29 stands in El Sohol area and Al Ta'akr mountain on axes 1 and 2.

In order to investigate the most significant factors affecting the grouping of the 29 stands, ANOVA (Analysis of Variance) was used to detect such factors. Soil moisture content and calcium content were the most significant factors in

determining different ecological groups with P value = 0.000, followed by chlorides with P value = 0.003 and organic carbon with P value = 0.014. (Table 3).

Table 3. Standard deviation (S. D.), Mean values and ANOVA P values of the soil variables in the studied sites.

Soil variable	TWINSPAN Group				P value
	A	B	C	D	
pH	7.5 ± 0.47	7.12 ± 0.17	7.12 ± 0.25	7.27 ± 0.95	0.594
EC	0.46 ± 0.26	0.26 ± 0.08	0.41 ± 0.31	0.35 ± 0.10	0.369
Cl	0.57 ± 0.49	0.10 ± 0.06	0.11 ± 0.07	0.10 ± 0.03	0.003
Mg	1.85 ± 1.31	1.48 ± 0.62	1.37 ± 0.45	1.51 ± 0.35	0.685
Ca	0.67 ± 0.21	0.42 ± 0.29	0.44 ± 0.14	1.07 ± 0.32	0.000
HCO ₃	109. ± 45.7	104. ± 33.1	97.6 ± 72.5	136. ± 131.	0.805
Carbonate	2.84 ± 0.51	1.34 ± 0.91	2.71 ± 0.73	3.98 ± 0.50	1.491
Organic Carbon	2.75 ± 0.58	2.76 ± 0.70	3.67 ± 0.69	2.01 ± 1.30	0.014
Moisture Content	16.3 ± 6.57	12.3 ± 4.32	9.80 ± 4.14	4.31 ± 3.32	0.000

DISCUSSION

The governorate of Ibbis named the Governorate of the Green Brigade and the reason for this is the dense vegetation cover on the mountains and the valleys. The climate of Ibb throughout the year, temperature is confined to between 27 to 33°C conditions suitable for growth as well as the high summer rainfall rate during the month of March to September makes the environmental conditions good for this dense vegetative growth.

A total of 111 species belonging to 51 families were recorded in 29 sites of the studied area. Family Asteraceae was the most abundant family comprising 12 species followed by Acanthaceae, Euphorbiaceae, Lamiaceae, Mimosaceae, Moraceae and Poaceae. Marei, (2006) reported that family Asteraceae is dominant family in south Sinai wadies.

The plant life form of the studied area shows that phanerophytes are the most dominant species representing 37%, followed by chamaephytes with 28 species representing 25%. Therophytes with 26 species representing 23%, cryptophytes with 13 species representing 11% and hemicryptophytes with only 2 species representing 1.8% of total species of the work.

These results supported by Aqlan, (2008).

On the other hand, Marei, et al (2016) studied Badan Mountain in Ibb governorate and recorded 71 species representing 38 families. Asteraceae was the prevalent family with 16 species followed by Poaceae (5 species) and Lamiaceae (4 species). Also the same authors studied life forms of species and mention that, chamaephytes was the most predominant life form with 33.8% of total species, followed by therophytes with 23.9%, Phanerophytes with 21.1%, cryptophytes with 19.7%, and hemicryptophytes with 1.4%.

Vegetation analysis of the study area showed that the trees and shrubs of the grassy plants support it (in the mountain of Al Ta'akr). seven sites and the herb of *Pennisetum setaceum* was observed only in one site (as dominant species) at plateau of AlTa'akr mountain.

In wadi El Ganat the dominated trees (*Ziziphus spina-christi* and *Acacia etbaica*) and shrubs of *Acacia mellifera* in 5 sites, while one site was dominated by *Arundo donax* represented the herbaceous case in the wadi.

In Najed El Bard the trees and shrubs also were dominated: 5 sites with *Ziziphus spina-christi*, *Acacia etbaica*, *Acacia mellifera* and *Acacia asak*

the shrubs were dominated in 3 sites (*Euphorbia cactus* and *Commiphora habessinica*), one site was dominated by the undershrub *Kleinia odora* and in one site the dominance was shared with both *Acacia etbaica* (tree) and *Commiphora habessinica* (shrub). In El Gabal El Akhdr 3 sites were dominated by shrubs (*Euphorbia cactus* and *Commiphora habessinica*), one site by undershrub (*Kleinia odora*) and one site by the herbaceous plant *Cynodon dactylon*. Also this species was present as co dominant in two sites in El Gabal El Akhader so the plateau and high slope of mountain attained their green color due to the coverage by *Cynodon dactylon*.

Al-Khulaidi & Pole (1989) mentioned that in the western high mountains of Yemen (including Ibb governorate), trees, shrubs and perennial herbs are dominated at the slope of mountain, terraces and even in the wadies. Plant communities of the present work are different from Al Ta'akr mountain than El Sohol area.

Community in plateau of Al Ta'akr mountain dominated by *Pennisetum setaceum*, this species was recorded by Wood (1997) who reported that this species grows well at elevation high as 3000m of sea level in plateau of mountain.

Communities in near plateau were dominated by *Rosa abyssinica* Wood (1997) reported that *R. abyssinica* is widespread on the escarpment and high plateau from 1500 to 3100 m above sea level. Al Khulidi (2000) recorded *R. abyssinica* on moderate steep slope mountains and hills, between 1900-2200m. Othman and El Naggar (2015) recorded *R. abyssinica* at high altitude mountains elevations above 1800 m high.

and communities of moderate and high slopes and also steep slopes were dominated by *Hypericum revolutum*. Wood (1997) recorded this species in Al Ta'akr mountain and reported that it grows above 2400 m and reaches 3200 m above sea level.

Communities in wadi bed are dominated by *Aruno donax*. Wood (1997) mention that this species is abundant along the margin of flowing wadies on the escarpment from 1500 to 1700 m. also may be found at 300 m. height.

Communities of the terraces were dominated

by *Ziziphus spina-christi* in wadi El Ganat and Najed El Bard, Wood (1997) reported that *Z. spina-christi* is found under 2500 m height of republic of Yemen and also this result is supported in the present work for all sites regarding the presence of this species below 2500 height.

Communities of slopes were dominated by *Acacia melifera* in wadi El Ganat and Najed El Bard Wood (1997) reported that *A. melifera* was found abundantly between 300 to 1700 m and not abundant at height more than 1900 m, Al Khulidi (2013) recorded this species on slopes north and north east Taiz (between 1200-1500m.) here this species forming a community type at height above 2000 m.

Acacia etbaica was recorded in wadi El Ganat and Najed El Bard Wood (1997) reported that *A. etbaica* is widespread between 1200 to 2300 m. Al Khulidi (2013) recorded *A. etbaica* on rocky wadi at plateau (1344 m). These woodlands are found on plains, plateaus, at the bottom of mountains and on moderate- steep slope mountains (between 1400-1800m.). Abdallah *et al.* (2016) recorded *A. etbaica* in El Baha at the altitude ranges from 1805 to 2020 m above sea level.

Communities dominated in the terraces in wadi El Ganat *Acacia asak* Wood (1997) reported that *A. asak* is abundant on steep well-drained mountain slope between 600 to 1700 m. Abdallah *et al.* (2016) recorded *A. asak* in El Baha at the altitude ranges from 1127 to 1756 m above sea level.

Communities of high and steep slopes were dominated by *Euphorbia cactus* Wood (1997) reported that *E. cactus* is widespread between 300 to 2000 m.

Commiphora habessinica Wood (1997) reported that *C. habessinica* is widespread between 1400 to 1900 m. And communities of plateau were dominated by *Ormocarpum dhofarense*, Wood (1997) recorded this species in Udayn (found in Ibb governorate) at 1500 m.

Kleinia odora Wood (1997) reported that *K. odora* is widespread between 600 to 2500 m. Abdallah *et al.* (2016) recorded *K. odora* in El Baha at the altitude ranges from 1805 to 2020 m above sea level.

Communities of moderate slope and mountain plateau were dominated by *Euphorbia inarticulate* Wood, (1997) reported that *E. inarticulate* is widespread between 300 to 2000 m.

Communities in high slopes were dominated by *Cynodon dactylon* Wood, (1997) reported that *C. dactylon* is one of the most widespread and abundant plant in Yemen occurring almost everywhere up to 2800 m. between 300 to 2000 m above sea level.

The application of TWINSpan on the vegetation data yielded four vegetation groups, 1- Group El Sohol area with height between 1800 to 2100 m including the following groups: Group A which is dominated by *Ziziphus spina-christi*, *Acacia etibca*, *Acacia milefera*, and *Arundo donxia*, Group B which is dominated by *Acacia etibca*, *Acacia origana*, *Ormocarpum dhofarense*, *Commiphora habessinica*, *Euphorbia inarticulate* and *Kleinia odora* and Group C which is dominated by *Acacia asak*, *Euphorbia cactus*, *Cadia purpurea*, *Acacia melifera*, *Cynodon dactylon*, and *Euphorbia inarticulate* as well as AlTa'akr mountain including one group (group D) with height between 2400 to 3000 m was dominated by *Rosa abyssinica*, *Hypericum revolutum* and *Pennisetum setaceum*. The results of Decorana and through the drawing sites of the top of AlTa'akr mountain related to axes 1 while the other sites are towards the direction of the axes 2 entirely and the results also showed the vast environmental difference between these groups of plants, while the other communities in Wadi El Ganat, El Gabal El Akhader and Najed El Bard to axes 2.

The results of analysis of variance showed that soil content of carbonate, calcium, moisture content and chlorides gave very high differences between the different groups, while the organic carbon content showed significant differences less than the previous elements, while PH, EC and HCO₃ showed no significant differences Fig.2

These Results were supported by Abd EL-Ghani & Marie (2006).

Carbonate and organic carbon showed significant differences between groups while supporting these results




Fig. 2. DCA ordination diagram for the TWINSpan groups of the 29 sites in in El Sohol area and AlTa'akr mountain on axes 1 and 2 in relation to different environmental factors.

Elevation play an important role in determine vegetation type, vegetation of AlTa'akr mountain (about 3000m above sea level) differ from El Sohol area (1800 to 2100 m above sea level) these area are near to each other being in the same governorate (Ibb governorate) but were different in vegetation characteristics and floristic composition of plants and communities. So, elevation is one of the factors determining phytogeographical regions and its communities in mountains, these results are in accordance with Lorphelin (1986) and Drever & Zobrist (1992).

RECOMMENDATION

This study, which was conducted on some plant communities in the province of Ibb.

1 - To strengthen the use of modern programs TWINSpan and DECORANA to facilitate understanding of the relationship between plant species and the environment.

2. In the future, GIS Geographic Information Systems (GIS) will be used to make accurate geographical maps and plant communities.

3- To limit and preserve plant species used in folk medicine, especially rare and threatened species.

4- Announcing the governorate of Ibb as anatural concerned area reserve in the future.

ACKNOWLEDGMENT

Thanks and appreciation to Mr. Esam Aqlan, assistant lecturer, Department of Biology, Fac-

ulty of Science, University of Ibb, for providing assistance in field trips during this study.

REFERENCES

- Abd El-Ghani, M. M. and Marie, A. H. (2006). Vegetation association of the endangered *Randonia africana* Coss. and its soil characteristics in an arid desert ecosystem of western Egypt. *Acta Bot. Croat.* 65 (1): 83-99.
- Abdullah, A., Al-Khulaidi, A., Akram, H. and Nageeb, A. (2016). Main vegetation types and plant species diversity along an altitudinal gradient of Al Baha region, Saudi Arabia. *Saudi Journal of Biological Sciences*. Vol. 23, 687 – 697.
- Al-Hubaishi, A. and Mueller-Hohenstein, K. (1984). *An Introduction to the Vegetation of Yemen: Ecological Basis, Floristic Composition, Human Influence*. Printed by as-Druck, D 6479 Schotten, Germany.
- Al Khulaidi and Pole, S. (1989). *Habitats of wild plants of the western part of the Republic of Yemen*. Ministry of Agriculture, Agriculture researches protection area, Republic of Yemen. (in Arabic).
- Al Khulaidi, A. A. (2000). *Flora of Yemen*. (a checklist). Sustainable Environmental Management Program (SEMP), YEM/97/100, sub-programm 11 and AREA, Sana'a, Yemen.
- Al Khulaidi, A. A. (2013). *Flora of Yemen. The Sustainable Natural Resource Management Project (SNRMP II)* EPA and UNDP.
- Aqlan, E. (2008). *Studies on the Flora of Ibb Governorate, Republic of Yemen*. Unpublished Thesis, Sana'a University, Yemen.
- Boulos, L. (1999). *Flora of Egypt. Vol 1, Azollaceae-Oxalidaceae*. Al Hadara Publishing, Cairo. 419pp.
- Boulos, L. (2000). *Flora of Egypt. Volume 2. Geraniaceae-Boraginaceae*. Al Hadara Publishing, Cairo. 352pp.
- Boulos, L. (2002). *Flora of Egypt. Volume 3. Verbenaceae-Compositae*. Al Hadara Publishing, Cairo. 373pp.
- Boulos, L. (2005). *Flora of Egypt. Volume 4. Monocotyledons*. Al Hadara Publishing, Cairo. 617pp.
- Boulos, L. (2009). *Flora of Egypt. Checklist. Revised Annotated Edition*. Al Hadara Publishing, Cairo. 410pp.
- Drever, J. I., and Zobrist, J. (1992). *Chemical Weathering of Silicate Rocks as A Function of Elevation in the Southern Swiss Alps*. *Geochimica et Cosmochimica Acta*, 56(8), 3209-3216.
- Forsskål, P. (1775). *Flora Aegyptiaco-Arabica*. Officina Mölleri, Hauniae.
- Jackson, M. L. (1967). *Soil Chemicals Analysis*. Prentice-Hall of India. Private, New Delhi, India.
- Lavranos, J. J. (1971a). Notes on the Succulent Flora of Northeast Africa and Southern Arabia, Part 1. *Cact. Succ. J. (U. S.)* 43 (1): 9-11.
- Lavranos, J. J. (1971b). Notes on the Succulent Flora of Northeast Africa and Southern Arabia, Part 2. *Cact. Succ. J. (U. S.)* 43 (2): 60-61.
- Lavranos, J. J. (1971c). *Senecio Delfersii* O. Schwartz; A Very Rare and Unusual Species from the Southern Yemen. *Cactus Succulent J.*
- Lorphelin, L. (1986). Weathering of Silt and Clay in Soils of A Toposequence in the Himalayas, Nepal. *Geoderma*, 39(2), 141-155.
- Marie, A. H. (2006). Floristic Composition and Vegetation Analysis of Downstream Parts of Two Wadies East of Al Qaa Plain, South Sinia. *Al Azaher J. Pharm. Sci.* Vol 34: 182 – 202.
- Marie, A. H., Elsaied, A. B. and Aqlan, E. C. (2016). *Phytosociological studies on Mount Badan, Ibb Governorate, Yemen Republic*. June, vol, 27 1 – 16.
- Mccune, B., and M. J. Mefford. (1999). *Pc-Ord for Windows Version 4.17. Multivariate Analysis of Ecological Data*. Mjmm Software, Glenenden Beach, Oregon, Usa.
- Miller, A. G. and Cope, T. A. (1996). *Flora of the Arabian Peninsula and Socatra*. (Vol. 1) Edinburgh Univ. Press. in Associ. Royal Botanic Garden. Eidenburgh, Kew, U.K.
- Othman, S. A. and El Naggar, S. M. (2015). *Vegetation patterns and floristic composition of Yemen*. *Current Life Sciences* 2015; 1 (3): 103-111.
- Piper, C. S. (1950). *Soil and Plant Analysis*. Univ. of Adelaide Press. Australia.
- Richards, L. A. (1954). *Diagnosis and Improvement of Saline and Alkaline Soils*. Usda Handbook No. 60, Washington, Dc.
- Santoso, S. (2014). *Spss 22 from Essential to Expert Skills*. Pt. Gramedia.
- Shukla, R. S. and Chandel, P. S. (1989). *Plant Ecology and Soil Science*. S. Chand & Co., New Delhi, India.
- Statistical Year Book (2001). *Central Statistical Organization, Ministry of Planning and Inter. Coop, Yemen*.
- Statistical Year Book (2002). *Central Statistical Organization, Ministry of Planning and Inter. Coop, Yemen*.
- Statistical Year Book (2003). *Central Statistical Organization, Ministry of Planning and Inter. Coop, Yemen*.
- Statistical Year Book (2004). *Central Statistical Organization, Ministry of Planning and Inter. Coop, Yemen*.
- Täckholm, V. (1974). *Students' Flora of Egypt*. Publ. Cairo Univ. Printing by Cooperative Printing Company Beirut, Pp 888.
- Ter Braak, C. J. F. (1988). *Canoco - A Fortran Program Or Canonical Community Ordination By (Partial) (Detrended) (Canonical) Correspondence Analysis, Principal Components Analysis and Redundancy Analysis, Version 2.1*. Wageningen, Agricultural Mathematics Group.
- Wood, J. R. I. (1997). *A Handbook of the Yemen Flora*. Royal Botanic Gardens, Kew, UK.

المجتمعات النباتية في جبل التعكر و منطقة السحول محافظة اب الجمهورية اليمنية

عبدة مرعى حامد مرعى

جامعة الازهر كلية العلوم قسم النبات و الميكروبيولوجي

الملخص

يقع جبل التعكر (3000 م) جنوب مدينه اب و منطقه السحول (1800 الي 2100 م) تقع شمال المدينه و الدراسه هي مقارنه بين تلك المنطقتان و المختلفتان من الناحيه التوبغرافيه تم تسجل 111 نوع نباتي و 51 فصيله نباتيه في 29 موقع في منطقه الدراسه و كانت الفصيله المركبه هي الفصيله الشائعه و السائده في منطقه الدراسه و الفصيله الاكاسيه و اللبنيه بعد ذلك و ضما 6 انواع نباتيه لكل منها ثم الفضائل الشفويه و التوتيه و النجيليه و احتوت كل فصيله منهم على 5 انواع نباتيه.

احتوى شكل الحياه Phanerophytes على 37% من الانواع المسجله ثم شكل الحياه chaemophytes احتوى على 25% من الانواع للمسجله ثم شكل الحياه Therophytes احتوى على 23% من الانواع المسجله ثم شكل الحياه Cryptophytes احتوى على 11% من الانواع المسجله و اخيرا شكل الحياه Hemicryptophytes احتوى على 1.8% من الانواع المسجله.

موقع واحد فقط سياده *Rosa abyssinica* و *Hypericum revolutum* في جبل التعكر وضح سياده الاشجار و الشجيرات في مصاطب *Ziziphus spina-christi* و ساد نبات *Arundo donax* للاعشاب المعمره في وادي الجنات ساد نبات في بطن الوادي في منحدر الوادي *Acacia melifera* و *Acacia etbaica* الوادي و نباتي

و في مصاطب الوادي و نبات *Acacia asak* في بطن الوادي و نبات *Ziziphus spina-christi* في نجد البرد ساد نبات في المنحدرات العاليه للوادي *Acacia melifera* في المنحدرات قليله الارتفاع و نبات *etbaica*

في المنحدرات القليله الارتفاع و نبات *Ormocarpum dhofarensis* في الجبل الاخضر ساد نبات في المنحدرات العاليه للجبل *Cynodon dactylon* في المنحدرات متوسطه الارتفاع و هضبه الجبل و كذلك نبات

تم استخدام برنامج TWINSpan في منطقه الدراسه لتقسيم المجموعات طبقا للانواع الدليله و كذلك برنامج DECORANA لتوضيح العلاقه بينتوزيع الانواع و العوامل البيئيه المختلفه و استخدم برنامج ANOVA لتوضيح عوامل التربيه التي على اساسها تم تقسيم الانواع في منطقه الدراسه اظهرت عوامل المحتوى الرطوبي و نسبه الكالسيوم في التربيه فروقا معنويه كبيره ثم الكلوريدات و كذلك الكربون العضوى