

6-1-2019

Section: Mathematics, Statistics, Computer Science, Physics and Astronomy

SOME UNUSUAL GENERATING INTEGRALS INVOLVING HERMITE POLYNOMIALS

Ahmed Youssef

Basic Science Department, Banha Faculty of Engineering, Banha University, Banha, Egypt., ahmed-yousef201480@yahoo.com

Follow this and additional works at: <https://absb.researchcommons.org/journal>



Part of the [Life Sciences Commons](#)

How to Cite This Article

Youssef, Ahmed (2019) "SOME UNUSUAL GENERATING INTEGRALS INVOLVING HERMITE POLYNOMIALS," *Al-Azhar Bulletin of Science*: Vol. 30: Iss. 1, Article 3.

DOI: <https://doi.org/10.21608/absb.2019.66074>

This Original Article is brought to you for free and open access by Al-Azhar Bulletin of Science. It has been accepted for inclusion in Al-Azhar Bulletin of Science by an authorized editor of Al-Azhar Bulletin of Science. For more information, please contact kh_Mekheimer@azhar.edu.eg.

SOME UNUSUAL GENERATING INTEGRALS INVOLVING HERMITE POLYNOMIALS

Ahmed. M. Youssef

Basic Science Department, Banha Faculty of Engineering, Banha University, Banha, Egypt.

Corresponding author: Ahmed-yousef201480@yahoo.com

ABSTRACT

In this paper we seek to find some bilateral generating functions for some orthogonal polynomials like Hermite, Charlier, Modified Laguerre and Gegenbauer polynomials. These bilateral generating functions can be obtained with the help of the unusual integral formula for Hermite polynomial seen in equation (1.1).

Keywords: Hermite polynomials, Modified Laguerre, Gegenbauer polynomials

1 INTRODUCTION

In a recent paper [1] the author established the following unusual integral for Hermite polynomials in the form

$$H_n(x) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) [2(x + iv)]^n dv \quad (1.1)$$

The aim of this paper is to find some unusual bilinear generating integrals by the help of the above equation (1.1).

Now, we can find the following bilinear generating functions of Hermite polynomials by using the unusual generating integral (1.1) as follows

$$\sum_{n=0}^{\infty} \frac{t^n}{2^n n!} H_n(x) H_n(y) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2 + 2xt(y + iv) - t^2(y + iv)^2) dv \quad (1.2)$$

The above bilinear generating relation (1.2) is deduced by Chatterjea S. K [1].

2. Bilateral Generating Function With Charlier and Hermite Polynomials

Here we notice the following linear generating relation involving Charlier polynomials [2]

$$\sum_{n=0}^{\infty} C_n(k; \alpha) \frac{t^n}{n!} = \exp(t) \left(1 - \frac{t}{\alpha}\right)^k \quad (2.1)$$

Now, we have

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{t^n}{n!} C_n(k; \alpha) H_n(x) \\
&= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left(\sum_{n=0}^{\infty} \frac{[2t(x+iv)]^n}{n!} C_n(k; \alpha) \right) dv \\
&= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \exp(2t(x+iv)) \left(1 - \frac{2t(x+iv)}{\alpha} \right)^k dv \\
&= \exp(2xt) \left(\frac{t}{\alpha} \right)^k \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2 + 2ivt) \left(\frac{\alpha}{t} - 2(x+iv) \right)^k dv
\end{aligned} \tag{2.2}$$

Now in order to evaluate (2.2) we shall prove the following integral formula

$$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2 + 2ivt) (2(x+iv))^k dv = \exp(-t^2) H_k(x-t) \tag{2.3}$$

In fact we notice that

$$\sum_{n=0}^{\infty} H_{n+k}(x) \frac{t^n}{n!} = \exp(2xt - t^2) H_k(x-t) \tag{2.4}$$

Now the left member of (2.4) can be written by virtue of (1.2) as follows

$$\begin{aligned}
& \frac{1}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{t^n}{n!} \int_{-\infty}^{\infty} \exp(-v^2) [2(x+iv)]^{n+k} dv \\
&= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \exp(2t(x+iv)) [2(x+iv)]^k dv \\
&= \exp(2xt) \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2 + 2ivt) [2(x+iv)]^k dv
\end{aligned}$$

Comparing the above integral with the right member of (2.4) we obtain (2.3).

Now we are in apposition to evaluate (2.2).

To this end we can alternatively write (2.3) in the form

$$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2 + 2ivt) (2(x-iv))^k dv = \exp(-t^2) H_k(x+t) \tag{2.5}$$

Thus it follows from (2.2) and (2.5) that

$$\sum_{n=0}^{\infty} C_n(k; \alpha) H_n(x) \frac{t^n}{n!} = \left(\frac{t}{\alpha}\right)^k \exp(2xt - t^2) H_k\left(\frac{2t^2 - 2xt + \alpha}{2t}\right) \quad (2.6)$$

3 Bilateral Generating Function with modified Laguerre and Hermite Polynomials

Here we notice the following linear generating relation involving modified Laguerre polynomials

$$\sum_{n=0}^{\infty} L_n^{k-n}(\alpha) t^n = \exp(-\alpha t) (1 + t)^k \quad (3.1)$$

Similarly by multiplying the left hand side of the previous equation by $H_n(x)$ and substitute from (1.2) we get

$$\begin{aligned} \sum_{n=0}^{\infty} L_n^{k-n}(\alpha) H_n(x) t^n &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left(\sum_{n=0}^{\infty} [2t(x + iv)]^n L_n^{k-n}(\alpha) \right) dv \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \exp(-2\alpha t(x + iv)) (1 + 2t(x + iv))^k dv \\ &= \frac{1}{\sqrt{\pi}} \exp(-2\alpha xt) \int_{-\infty}^{\infty} \exp(-v^2 - 2\alpha tiv) (1 + 2t(x + iv))^k dv \end{aligned}$$

By replacing (t) by $(-t)$ in the previous equation and substitute from (2.3) we get

$$\begin{aligned} \sum_{n=0}^{\infty} L_n^{k-n}(\alpha) H_n(x) (-t)^n &= \frac{1}{\sqrt{\pi}} \exp(2\alpha xt) \int_{-\infty}^{\infty} \exp(-v^2 + 2\alpha tiv) (1 - 2t(x + iv))^k dv \\ &= \frac{1}{\sqrt{\pi}} \exp(2\alpha xt) (t)^k \int_{-\infty}^{\infty} \exp(-v^2 + 2\alpha tiv) \left(\frac{1}{t} - 2(x + iv)\right)^k dv \\ &= \frac{1}{\sqrt{\pi}} \exp(2\alpha xt) (t)^k \int_{-\infty}^{\infty} \exp(-v^2 + 2\alpha tiv) \left(2\left(\frac{1}{2t} - x - iv\right)\right)^k dv \\ &= \exp(2\alpha xt) (t)^k \exp(-\alpha^2 t^2) H_k\left(\frac{1}{2t} - x + \alpha t\right) \\ &= t^k \exp(2\alpha xt - \alpha^2 t^2) H_k\left(\frac{2\alpha t^2 - 2xt + 1}{2t}\right) \end{aligned}$$

Hence;

$$\begin{aligned} & \sum_{n=0}^{\infty} L_n^{k-n}(\alpha) H_n(x) (-t)^n \\ &= t^k \exp(2\alpha xt - \alpha^2 t^2) H_k\left(\frac{2\alpha t^2 - 2xt + 1}{2t}\right) \end{aligned} \quad (3.2)$$

It may be of interest to note that we can obtain equation (3.2) from equation (2.5) so we have to prove that:

$$C_n(k, \alpha) = (-\alpha)^{-n} n! L_n^{k-n}(\alpha) \quad (3.3)$$

From Laguerre polynomial $L_n^x(\alpha)$

$$L_n^x(\alpha) = \sum_{r=0}^n \frac{(-1)^r (1+x)_n}{r! (n-r)! (1+x)_r} \alpha^r$$

Let $x = k - n$, then

$$\begin{aligned} L_n^{k-n}(\alpha) &= \sum_{r=0}^n \frac{(-1)^r (1+k-n)_n}{r! (n-r)! (1+k-n)_r} \alpha^r \\ &= \sum_{r=0}^n \frac{(-1)^r (1+k-n)_n}{r! (n-r)! (1+k-n)_r} \alpha^r \\ &= \sum_{r=0}^n \frac{(-1)^r (-k)_{n-r} (-1)^{n-r}}{r! (n-r)!} \alpha^r \\ &= \sum_{r=0}^n \frac{(-1)^n (-k)_{n-r}}{r! (n-r)!} \alpha^r \end{aligned}$$

Multiplying both sides of the previous equation by $n! (-\alpha)^{-n}$

$$\begin{aligned} (-\alpha)^{-n} n! L_n^{k-n}(\alpha) &= \sum_{r=0}^n \frac{(-1)^n (-k)_{n-r}}{r! (n-r)!} \alpha^r (-\alpha)^{-n} n! \\ &= \sum_{r=0}^n \frac{n! (-k)_{n-r}}{r! (n-r)!} \alpha^{r-n} \end{aligned}$$

Let $m = n - r$, hence

$$\begin{aligned} &= \sum_{m=0}^n \frac{n! (-k)_m}{m! (n-m)!} \alpha^{-m} \\ &= \sum_{m=0}^n \binom{n}{m} (-k)_m \alpha^{-m} \end{aligned}$$

Since $(-k)_m = (-1)^m \frac{k!}{(k-m)!}$ hence;

$$\begin{aligned} (-\alpha)^{-n} n! L_n^{k-n}(\alpha) &= \sum_{m=0}^n (-1)^m \binom{n}{m} \frac{k!}{(k-m)!} \alpha^{-m} \\ &= \sum_{m=0}^n (-1)^m \binom{n}{m} \binom{k}{m} \frac{m!}{(\alpha)^m} \\ &= C_n(k, \alpha) \end{aligned}$$

Hence;

$$C_n(k, \alpha) \frac{\alpha^n}{n!} = (-1)^n L_n^{k-n}(\alpha) \quad (3.4)$$

Finally by substitute from equation (3.4) in equation (2.5) we get

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{t^n}{n!} (-1)^n L_n^{k-n}(\alpha) \frac{n!}{\alpha^n} H_n(x) \\ = \left(\frac{t}{\alpha}\right)^k \exp(2xt - t^2) H_k\left(\frac{2t^2 - 2xt + \alpha}{2t}\right) \end{aligned}$$

Hence;

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{-t}{\alpha}\right)^n L_n^{k-n}(\alpha) H_n(x) \\ = \left(\frac{t}{\alpha}\right)^k \exp(2xt - t^2) H_k\left(\frac{2t^2 - 2xt + \alpha}{2t}\right) \end{aligned} \quad (3.5)$$

We notice that we can obtain the same result (3.5) from (3.2) by replacing (t) by (t/α) .

4 Bilateral Generating Function With Gegenbaur and Hermite polynomials

Here we notice the following linear generating relation involving Gegenbaur polynomials

$$\sum_{n=0}^{\infty} C_n^v(x) t^n = (1 - 2xt + t^2)^{-v} \quad (4.1)$$

Now by multiplying the left hand side of (3.1) by $H_n(x)$ and substituting by (1.2) we get (3.2)

$$\begin{aligned} \sum_{n=0}^{\infty} C_n^v(x) t^n H_n(x) &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left(\sum_{n=0}^{\infty} [2t(x + iv)]^n C_n^v(x) \right) dv \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left[1 - 4xt(x + iv) + (2t(x + iv))^2 \right]^{-v} dv \end{aligned} \quad (4.2)$$

In particular case, if we take $x = 1$ in the above equation (4.1), then we have

$$\begin{aligned} \sum_{n=0}^{\infty} C_n^v(1) t^n &= (1 - 2t + t^2)^{-v} \\ &= (1 - t)^{-2v} \\ &= \sum_{n=0}^{\infty} \frac{(2v)_n}{n!} t^n \end{aligned}$$

Hence;

$$C_n^v(1) = \frac{(2v)_n}{n!}$$

Secondly substitute in (4.2) we get

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(2v)_n}{n!} H_n(x) t^n &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left[1 - 4t(x + iv) + (2t(x + iv))^2 \right]^{-v} dv \end{aligned}$$

Finally we get;

$$\begin{aligned} & \sum_{n=0}^{\infty} (2v)_n e^{2xt-t^2} \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(-v^2) \left[1 - 4t(x + iv) + (2t(x + iv))^2 \right]^{-v} dv \quad (4.3) \end{aligned}$$

REFERENCES

- [1] Chatterjea, S. K., My favourite proof of Mehler's formula, to appear in Jour. Pure math. Vol. 2 (1982).
- [2] Lee, P. A., Some generating function involving the Charlier's and Hermite polynomials, Nanta. Math. 8 (1975), 83-87.
- [3] Rainville, E. D., Special function, New Yourk (1960).

الملخص العربي

يتناول هذا البحث بعض التعاريف الهامة بالنسبة الي كثيرة حدود هارميت بالاضافة الي عملية الاندماج مع بعض كثيرة الحدود الاخرى مثال ذلك كثيرة حدود لاجير وكثيرة حدود تشارلز وكذلك الاندماج مع كثيرة حدود جاجن بيور مما يعطي اضافة للبحث في مجال تعريف الدوال الخطية والدوال ثنائي الخطية .